

A.R. Clapham

Downing

Idestrichum fennoscandicum

Desmanella

Desmanella

Hypnum cupressinum

to var. *filiforme*

Hylacomium luguberrimum

Thuidium

Brachythecium

Mnium und.

Catharinae

A. R. Clapham

Downing

GENERA OF BRITISH
PLANTS

CAMBRIDGE UNIVERSITY PRESS

London: FETTER LANE, E.C.

C. F. CLAY, MANAGER



Edinburgh: 100, PRINCES STREET

London: WILLIAM WESLEY AND SON, 28, ESSEX STREET, STRAND

Berlin: A. ASHER AND CO.

Leipzig: F. A. BROCKHAUS

New York: G. P. PUTNAM'S SONS

Bombay and Calcutta: MACMILLAN AND CO., LTD.

All rights reserved

GENERA OF BRITISH PLANTS

ARRANGED ACCORDING TO
ENGLER'S *SYLLABUS DER PFLANZENFAMILIEN*
(SEVENTH EDITION 1912)

WITH THE ADDITION OF THE CHARACTERS
OF THE GENERA

BY

HUMPHREY G. CARTER, M.B., CH.B.

Cambridge:
at the University Press

1913

GENERAL OF BRITISH PLANTS

THE UNIVERSITY PRESS

PRINTED BY JOHN CLAY, M.A.

AT THE UNIVERSITY PRESS

WITH THE ADDITION OF THE CHARACTERS
OF THE FLORA

Cambridge:

PRINTED BY JOHN CLAY, M.A.

AT THE UNIVERSITY PRESS

THE UNIVERSITY PRESS

Cambridge:

at the University Press

1872

PREFACE

WHAT is known as Engler's System of the Classification of Plants embodied in the *Natürliche Pflanzenfamilien* and in the successive editions of the *Syllabus der Pflanzenfamilien*, in the opinion of many modern systematic botanists represents, as a whole, the nearest approach to a natural system that we possess.

While specialists may criticise it in one detail or another, and while it is certain that we are as yet far from finality, owing to the numerous lacunae which still exist in our knowledge of different groups, there can be little if any doubt that no other published system of classification rivals Professor Engler's in broad grasp of principle and masterly treatment of complicated detail. It is the conspicuous merit of Professor Engler and his collaborators that they have kept constantly presented to botanists in the successive editions of the *Syllabus* the best scheme that could in their judgment be devised at the moment.

The present work is intended to familiarise students of British Vascular Plants with Engler's system in its latest form and thus both to habituate British floristic students to the use of a more natural system than that

to which they have been accustomed in the British Floras that have hitherto appeared, and to serve as an introduction to the use of the *Syllabus*, the *Natürliche Pflanzenfamilien* and the *Pflanzenreich*.

The major divisions of the vascular plants with their diagnoses are taken, to a large extent, *verbatim* from the *Syllabus der Pflanzenfamilien* (7th Edition 1912).

Professor Engler and his publishers have most kindly and readily accorded their assent to this very free use of their publication.

In a few cases Engler's arrangement has been departed from.

Thus the nomenclature of the larger divisions of Pteridophyta has been somewhat modified in order to secure uniformity in the terminations of the names of Orders (Reihen). Prof. Bower's arrangement of the Leptosporangiate Ferns into Simplicis Mixtae and Gradatae has been adopted as representing the most rational grouping, though the distinct suborder of Filicales, Hydropteridineae, has been retained owing to the difficulty of assigning with certainty the Marsiliaceae and Salviniaceae to their proper places in the Eufilianeae series.

The Family Polypodiaceae, practically equivalent to the Mixtae of Bower, has been retained and divided into tribes according to the arrangement of the *Syllabus*, since our knowledge of the Systematics of this intricate

group is not sufficiently advanced to admit of a grouping of the Mixtae into Natural Families. In dealing with the British Genera alone this arrangement does not appear to do much violence to natural affinities so far as they have been ascertained.

Among the Flowering Plants Warming's arrangement of the Urticales has been followed mainly because of the difficulty of defining the Family Moraceae so as to include Cannabis and Humulus (subfamily Cannaboideae of Engler).

In all the other Orders the arrangement of the *Syllabus* has been closely followed, though the actual diagnoses, especially in cases where an Order or Family is poorly represented in Britain, have been to some extent modified.

The Characters of the Genera have either been obtained from various sources, such as Hooker's *Student's Flora* and Garcke's *Flora von Deutschland*, or are original, and they will, it is hoped, prove useful to students.

The following books, besides the *Syllabus* and *Pflanzenfamilien*, have been constantly used by me during the compilation of this work.

Ascherson and Graebner. *Flora des Nordostdeutschen Flachlandes*, 1898-9.

Blytt. *Haandbog i Norges Flora*, 1906.

Garcke. *Flora von Deutschland*. Edited by Dr Franz Niedenzu, 1908.

Hooker, Sir J. D. *The Student's Flora of the British Islands*, 1884.

Rendle. *The Classification of Flowering Plants*. Vol. I.

Warming. *Frøplanterne*, 1912.

Willis, J. C. *Flowering Plants and Ferns*, 1908.

My best thanks are due to Mr Tansley for help and advice with the Pteridophytes, to Professor Seward for reading the proofs of the Pteridophytes and Gymnosperms, and to Dr Moss who has read the entire proofs and given me much valuable advice and assistance throughout.

H. G. C.

CAMBRIDGE,

August 1913.

SYNOPTIC INDEX

The numbers in the first column refer to pages in this book, those in the second column refer to pages in Engler's Syllabus (Ed. 7, 1912).

Division 1. EMBRYOPHYTA ASIPHONOGAMA (ARCHEGONIATAE)	1	74
[Subdivision 1. Bryophyta]	1	75
Subdivision 2. PTERIDOPHYTA	1	88
Class 1. EQUISETARIAE (Equisetales in Syllabus)	1	97
Order Equisetales (Euequisetales in Syllabus)	2	97
Fam. EQUISETACEAE	2	97
Class 2. LYCOPODIARIAE (Lycopodiales in Syllabus)	2	98
Order 1. Lycopodiales (Lycopodiales eligulatae and ligulatae of Syllabus) .	2	98-99
Fam. 1. LYCOPODIACEAE	2	98
Fam. 2. SELAGINELLACEAE	3	99
Order 2. Isoëtales (Class in Syllabus) .	3	101
Fam. ISOËTACEAE	3	101
Class 3. FILICARIAE (Filicales in Syllabus).	3	89

Order 1.	Filicales (Filicales Leptosporan-		
	giatae in Syllabus)	4	89
Fam. 1.	OSMUNDACEAE	4	94
Fam. 2.	HYMENOPHYLLACEAE	5	90
Fam. 3.	POLYPODIACEAE	5	90
Fam. 4.	MARSILIACEAE	7	95
Fam. 5.	SALVINIACEAE	7	95
Order 2.	Ophioglossales	7	96
Fam.	OPHIOGLOSSACEAE	8	97
Division 2.	EMBRYOPHYTA SIPHO-		
	NOGAMA (<i>Phanerogamae</i> auct.)	8	102
Subdivision 1.	GYMNOSPERMAE	8	103
Class 1.	CONIFERAE	9	106
Fam. 1.	TAXACEAE	9	106
Fam. 2.	PINACEAE	9	108
Subdivision 2.	ANGIOSPERMAE	10	113
Class 1.	MONOCOTYLEDONES	10	117
A.	Orders with preponderating incon-		
	stancy in the number of floral		
	leaves	10	117
a.	<i>Typical achlamydeous (not apopetalous) flowers</i>		
	<i>still occur</i>	10	117
a.	<i>Achlamydeous flowers predominate. Great incon-</i>		
	<i>stancy in number of stamens and also</i>		
	<i>of carpels</i>	10	117
Order 1.	Pandanales	10	117
Fam. 1.	TYPHACEAE	11	118
Fam. 2.	SPARGANIACEAE	11	118
β.	<i>Naked flowers still present: but in this order there</i>		
	<i>occur all possible gradations from A-</i>		
	<i>chlamydy to Heterochlamydy, also from</i>		
	<i>Hypogyny to Epigyny. Sporophylls defi-</i>		
	<i>nite or indefinite</i>	11	118

Order 2.	Alismatales	11	118
Fam. 1.	POTAMOGETONACEAE	11	119
Fam. 2.	NAIADACEAE	12	120
Fam. 3.	*APONOGETONACEAE	12	120
Fam. 4.	SCHEUCHZERIAACEAE	13	120
Fam. 5.	ALISMATACEAE	13	120
Fam. 6.	BUTOMACEAE	14	121
Fam. 7.	HYDROCHARITACEAE	14	121
γ.	<i>Naked flowers dominant. Stamens rarely, and carpels never, indefinite</i>	15	123
Order 3.	Glumiflorae	15	123
Fam. 1.	GRAMINACEAE	15	123
Fam. 2.	CYPERACEAE	20	127
b.	<i>Achlamydeous flowers rare: when they occur they are mostly due to reduction and are correlated with spathe-development. Definite number of stamens and carpels dominant, but numerous stamens and more than three carpels often occur'</i>	21	128
Order 4.	Spathiflorae	22	133
Fam. 1.	ARACEAE	22	133
Fam. 2.	LEMNACEAE	23	136
B.	Orders with typically pentacyclie flowers. Whorls typically isomerous, mostly 3-merous	23	137
a.	<i>Flowers homochlamydeous to heterochlamydeous, very rarely by reduction naked. Bracteoid perianths still occur but petaloid perianths predominate. Hypogyny and Actinomorphy preponderate</i>	23	137
Order 5.	Farinosae	23	137
Fam.	ERIOCAULACEAE	23	138

Order 6. Liliiflorae	24	141
Fam. 1. JUNCACEAE	24	141
Fam. 2. LILIACEAE	24	141
Fam. 3. AMARYLLIDACEAE	27	146
Fam. 4. DIOSCOREACEAE	28	147
Fam. 5. IRIDACEAE	28	148
<i>b. Flowers homochlamydeous to heterochlamydeous but the perianth is always petaloid. Epigyny throughout. Zygomorphy dominant</i>							
Order 7. Microspermae	29	151
Fam. ORCHIDACEAE	30	152
Class 2. DICOTYLEDONES	33	156
Subclass 1. ARCHICHLAMYDEAE	33	156
<i>a. Amentiflorae (Moss). Mostly trees or shrubs. ♂ flowers usually, and ♀ flowers often, in catkins. Perianth absent or haplo- chlamydeous and bracteoid.</i>							
Order 1. Salicales	33	157
Fam. SALICACEAE	33	158
Order 2. Myricales	34	159
Fam. MYRICACEAE	34	159
Order 3. *Juglandales	34	160
Fam. *JUGLANDACEAE	34	160
Order 4. Fagales	34	161
Fam. 1. BETULACEAE	35	162
Fam. 2. FAGACEAE	35	162
Order 5. Urticales	36	164
Fam. 1. ULMACEAE	36	164
Fam. 2. CANNABACEAE (<i>Moraceae, Can- naboideae in Syllabus</i>)	36	167
Fam. 3. URTICACEAE	37	167

<i>b.</i>	Petaloidae (Moss). Flowers haplo-		
	chlamydeous, perianth often		
	petaloid. Diplochlamydeous		
	flowers rare	37	168
Order 6	Santalales	37	170
	Fam. 1. SANTALACEAE	38	170
	Fam. 2. LORANTHACEAE	38	172
Order 7.	Aristolochiales	38	176
	Fam. ARISTOLOCHIACEAE	38	176
Order 8.	Polygonales	39	177
	Fam. POLYGONACEAE	39	177
<i>c.</i>	(Consisting of the one order Centro-		
	spermae.) Corolla usually		
	haplochlamydeous, sepaloid		
	or petaloid. Heterochlamydy,		
	however, is not uncommon	40	179
Order 9.	Centrospermae	40	179
	Fam. 1. CHENOPODIACEAE	40	179
	Fam. 2. *AMARANTACEAE	41	181
	Fam. 3. *AIZOACEAE	42	183
	Fam. 4. PORTULACACEAE	42	183
	Fam. 5. CARYOPHYLLACEAE	43	184
<i>d.</i>	Heterochlamydeae (Moss). Orders with		
	predominant heterochlamydy	45	185
<i>a.</i>	<i>Apocarpy and Hypogyny predominant</i>	45	185
Order 10.	Ranales	45	185
	Fam. 1. NYMPHAEACEAE	45	185
	Fam. 2. CERATOPHYLLACEAE	46	186
	Fam. 3. RANUNCULACEAE	46	187
	Fam. 4. BERBERIDACEAE	48	188
<i>β.</i>	<i>Syncarpy and Hypogyny predominant</i>	48	195
Order 11.	Rhoeadales	49	195
	Fam. 1. PAPAVERACEAE	49	195
	Fam. 2. CRUCIFERAE	50	197
	Fam. 3. RESEDACEAE	54	201

Order 12.	Sarraceniales	54	201
	Fam. DROSERACEAE	54	202
γ.	<i>Apocarpy and Hypogyny still occur but Perigyny becomes more frequent. By sinking of the Gynaecium into the hollow flower-axis Syncarpy and Epigynous insertion of the perianth and stamens also takes place</i>					54	203
Order 13.	Rosales	54	203
	Fam. 1. CRASSULACEAE	55	204
	Fam. 2. SAXIFRAGACEAE	55	204
	Fam. 3. *PLATANACEAE	56	209
	Fam. 4. ROSACEAE	57	210
	Fam. 5. LEGUMINOSAE	59	215
δ.	<i>The flowers have predominantly 5 or 4 whorls. Apocarpy and Isomery still occur, but Syncarpy and Oligomery of the Gynaecium preponderate. Pleiomery of Gynaecium rare</i>					61	223
Order 14.	Geraniales	61	223
	Fam. 1. GERANIACEAE	62	223
	Fam. 2. OXALIDACEAE	63	224
	Fam. 3. LINACEAE	63	226
	Fam. 4. POLYGALACEAE	63	235
	Fam. 5. EUPHORBIACEAE	63	236
	Fam. 6. CALLITRICHACEAE	64	241
Order 15.	Sapindales	64	241
	Fam. 1. BUXACEAE	65	241
	Fam. 2. EMPETRACEAE	65	241
	Fam. 3. AQUIFOLIACEAE	65	244
	Fam. 4. CELASTRACEAE	65	245
	Fam. 5. *STAPHYLEACEAE	66	246
	Fam. 6. ACERACEAE	66	246
	Fam. 7. *HIPPOCASTANACEAE	66	247
	Fam. 8. BALSAMINACEAE	67	249

Order 16.	Rhamnales	67	250	
	Fam. RHAMNACEAE	67	250	
Order 17.	Malvales	67	252	
	Fam. 1. TILIACEAE	68	253	
	Fam. 2. MALVACEAE	68	253	
<i>ε. Flowers spirocyclic or of 5-4 whorls. Syncarpy the rule (Apocarpy in the more primitive, exotic families). Tendency for gynaeceium to become sunk in flower-axis</i>							69	257
Order 18.	Parietales	69	257	
	Fam. 1. GUTTIFERAE	69	261	
	Fam. 2. ELATINACEAE	70	264	
	Fam. 3. FRANKENIACEAE	70	264	
	Fam. 4. *TAMARICACEAE	70	264	
	Fam. 5. CISTACEAE	71	266	
	Fam. 6. VIOLACEAE	71	266	
<i>§. The flowers are cyclic and the sinking of the gynaeceium into the hollow flower-axis is general: connation of gynaeceium and flower-axis predominates</i>							71	274
Order 19.	Myrtiflorae	71	274	
	Fam. 1. THYMELAEACEAE	72	275	
	Fam. 2. ELAEAGNACEAE	72	276	
	Fam. 3. LYTHRACEAE	72	276	
	Fam. 4. EPILOBIACEAE (Oenotheraceae)	73	283	
	Fam. 5. HALORRHAGACEAE	74	284	
	Fam. 6. HIPPURIDACEAE	74	285	
Order 20.	Umbellales	74	285	
	Fam. 1. ARALIACEAE	74	285	
	Fam. 2. UMBELLIFERAE	75	286	
	Fam. 3. CORNACEAE	80	290	
Subclass 2.	METACHLAMYDEAE	80	291	
A. Polypetaly, as well as sympetaly, occurs. Two whorls or one								

	whorl of stamens. Hypogyny predominates, but epigyny also occurs	80	291
Order 1.	Ericales	80	291
	Fam. 1. PIROLACEAE	80	291
	Fam. 2. ERICACEAE	81	292
Order 2.	Primulales	83	295
	Fam. PRIMULACEAE	83	295
Order 3.	Plumbaginales	84	296
	Fam. PLUMBAGINACEAE	85	296
B. (C. in Syllabus).	Sympetaly dominant. Stamens always in one whorl. Union of Carpels sometimes imperfect. Hypogyny usual .	85	300
Order 4.	Contortae	85	300
	Fam. 1. OLEACEAE	85	300
	Fam. 2. GENTIANACEAE	86	303
	Fam. 3. APOCYNACEAE	87	303
C. (D. in Syllabus).	Sympetaly constant. Only one whorl of stamens present. Carpels usually 2, always perfectly united. Zygomorphy frequent .	87	308
a.	<i>Perianth hypogynous</i>	87	308
Order 5.	Tubiflorae	88	308
	Fam. 1. CONVULVULACEAE	88	308
	Fam. 2. POLEMONIACEAE	88	310
	Fam. 3. BORRAGINACEAE	89	311
	Fam. 4. VERBENACEAE	91	313
	Fam. 5. LABIATAE	91	315
	Fam. 6. SOLANACEAE	94	320
	Fam. 7. SCROPHULARIACEAE	95	322
	Fam. 8. OROBANCHACEAE	97	327
	Fam. 9. LENTIBULARIACEAE	97	329

Order 6. Plantaginales	98	332
Fam. PLANTAGINACEAE	98	332
<i>b. Perianth epigynous</i>	98	333
<i>a. Stamens free</i>	98	333
Order 7. Rubiales	99	333
Fam. 1. RUBIACEAE	99	333
Fam. 2. CAPRIFOLIACEAE	99	336
Fam. 3. ADOXACEAE	100	337
Fam. 4. VALERIANACEAE	101	337
Fam. 5. DIPSACACEAE	101	338
<i>β. Stamens conniving or partly united</i>	101	338
Order 8. Cucurbitales	102	338
Fam. CUCURBITACEAE	102	338
Order 9. Campanulatae	102	340
Fam. 1. CAMPANULACEAE	102	340
Fam. 2. COMPOSITAE	103	342

GLOSSARY OF WORDS RELATING TO PERIANTH

Achlamydeous.	With no perianth (naked).
Apochlamydeous.	Perianth absent by reduction.
Apopetalous.	Corolla absent by reduction.
Haplochlamydeous.	Having a perianth consisting of a single whorl.
Diplochlamydeous.	Having a perianth consisting of two whorls.
Homochlamydeous.	Diplochlamydeous with both whorls of the same kind.
Heterochlamydeous.	Diplochlamydeous with whorls clearly unlike each other.

In Floral Formulae

K = Calyx. C = Corolla. A = Androecium. G = Gynaecium.

* = not native.

⊙ = doubtfully native.

Division 1. EMBRYOPHYTA ASIPHONOGAMA (ARCHEGONIATAE).

Plants with marked alternation of generations. *Sexual generation* (*gametophyte*) bearing *antheridia*, in which *spermatozoids* arise, and *archegonia*, each containing an *egg-cell*. The fertilized egg-cell gives rise to an *embryo*, which for some time or throughout life remains attached to, and nutritively dependent on, the gametophyte. The embryo grows into the asexual generation (*sporophyte*), eventually producing *spores*, which again give rise to the gametophytes.

[Subdivision 1. Bryophyta (not included in this work). Gametophyte thalloid or leafy, free living. Sporophyte (sporogonium) leafless, attached throughout life to the gametophyte : no true roots.]

Subdivision 2. PTERIDOPHYTA. *Gametophyte* (*prothallus*) thalloid, usually vegetating independently, rarely completely enclosed in the spore. *Sporophyte* a cormophytic plant with true leaves and roots, early becoming independent of the gametophyte, and bearing *spores* in separate organs (*sporangia*) borne on or in connection with the leaves.

Class 1. EQUISETARIAE. *Leaves* generally small in relation to the stem, in alternating whorls. *Stele* with separate collateral bundles. Peltate *sporangiphores* aggregated in cones. *Sporangia* bulky, attached to the

inner surface of the lamina of the sporangiophore. Homosporous or (in some extinct forms) heterosporous.

Order **Equisetales** (including all living forms). Homosporous.

Family EQUISETACEAE. *Rhizomes* subterranean often bearing distinct sterile and fertile shoots. Bases of leaf-whorls united into sheaths enclosing the base of the internode above. *Branching* axillary. *Epispore* splitting into two spirally wound threads (*elaters*). *Prothalli* branched, often functionally dioecious. *Antheridia* and *archegonia* sunken. *Spermatozoids* multiciliate.

Only genus: **Equisetum.**

Class 2. **LYCOPODIARIAE.** *Leaves* usually small in relation to stem. *Stem* with one or more haplosteles†. *Sporangia* single in axils, or on adaxial faces of fertile leaves. Homosporous or heterosporous. *Spermatozoids* biciliate.

Order 1. **Lycopodiales.** Small leaved herbs of erect or creeping habit. *Sporophylls* mostly aggregated in *cones*.

Suborder 1. **ELIGULATAE.** Homosporous forms without *ligule*.

Family LYCOPODIACEAE. Erect or suberect herbs with modified *haplostele*. *Prothalli* generally subterranean, saprophytic, with symbiotic fungus. *Embryo* often living a subterranean life for some years.

Only British genus: **Lycopodium.**

† Haplostele. The cauline vascular cylinder in its simpler form, in which the centre is occupied by xylem, and this is typically surrounded by phloem, pericycle and endodermis (Brebner, *Ann. Bot.* 1902, p. 523 and Tansley, *Evolution of Filicinean Vascular System*, 1908).

Suborder 2. **LIGULATAE**. Heterosporous forms with ligule.

Family **SELAGINELLACEAE**. Erect or suberect herbs. Haplostelic or polystelic. *Rhizophores* often arising exogenously and bearing endogenous roots. *Leaves* rarely spiral, more often in two dorsal and two ventral rows. *Sporophylls* aggregated in cones, which contain both macro- and microsporophylls. *Macrosporangia* typically with four macrospores, *microsporangia* with numerous microspores. *Embryo* with two cotyledons.

Only genus: **Selaginella**.

(The British species is haplostelic with spirally arranged leaves.)

Order 2. **Isoëtales**. Submerged subaquatic or terrestrial plants with very short unbranched fleshy stems, bearing alternating whorls of sterile and fertile, elongated, subulate or filiform leaves. *Stele* with anomalous secondary thickening. *Sporangia* bulky, single, in pit on adaxial face of leaf base adjoining the ligule. *Macrosporangia* with numerous macrospores, *microsporangia* with very numerous microspores. Sterile *trabeculae* between the spores. *Spermatozoids* multiciliate.

Only family: **ISOËTACEAE**.

Only genus: **Isoëtes**.

Class 3. **FILICARIAE**. *Leaves* almost always large in relation to the stem, often highly compound, mostly with circinate vernation. *Stem* haplostelic, solenostelic†

† Solenostele. A stele in which the vascular tissue is arranged in a hollow cylinder of xylem lined within and without by phloem, pericycle and endodermis, and the continuity of which is interrupted by leaf-gaps which do not overlap (Tansley, *l.c.*).

or dictyostelic†, departure of leaf-traces in all but haplostelic forms making definite *leaf-gap* in the stele. *Sporangia* on the undersides or on the edges of ordinary foliage leaves, or of modified leaves (*sporophylls*) with reduced laminae; in the latter case the sporophylls are not aggregated on special shoots, or special regions of the shoot. *Spermatozoids* multiciliate.

Order 1. **Filicales**. *Sporangia* mostly arising from single cells, more rarely from groups or from several layers of cells, usually in definite aggregates (*sori*). *Prothallus* not subterranean, generally flat.

Suborder 1. FILICINEAE. Wall of the sporangium typically with a prominent cell row or cell group (*annulus*) concerned with the dehiscence of the sporangium. *Spores* all alike producing comparatively large free-living monoclinal or diclinal prothalli.

SIMPLICES (Bower). *Sporangia* grouped in sori each containing few sporangia of simultaneous origin on a flat or nearly flat receptacle or spread over the lower surface of the fertile leaf, relatively large, sessile or subsessile, with a large spore output. *Annulus* transverse or oblique or (as in the British genus) represented by a group of thickened cells. *Indusium* typically absent.

Family OSMUNDACEAE. *Stem* upright covered with the bases of spirally arranged leaves. *Stele* with continuous mantle of peripheral *phloem* and anastomosing *xylem strands* surrounding a *pith*. *Petiolar vascular*

† Dictyostele. A stele in which the hollow cylinder of the solenostelic type is interrupted by leaf-gaps which overlap (Brebner, *l.c.*).

strand horseshoe-shaped in cross section. *Sporangia* massive, shortly stalked with vertical dehiscence, not arranged in *sori*, but thickly set on the surface of special, often reduced, fronds or pinnae.

Only British genus :

Osmunda.

GRADATAE (Bower). *Sporangia* basipetal, on a more or less elongated receptacle, of moderate size, shortly stalked, with medium spore output. *Annulus* oblique. *Indusium* a basal cup, entire or 2-lipped.

Family HYMENOPHYLLACEAE. *Rhizomes* creeping, haplostelic, bearing fronds with thin filmy laminae which have no stomata or intercellular spaces in the mesophyll. *Sori* marginal on the ends of excurrent veins.

Prothalli filiform. *Indusium* tubular. **Trichomanes.**

Prothalli flat. *Indusium* 2-lipped.

Hymenophyllum.

MIXTAE (Bower). *Sporangia* of various ages mixed without regular orientation, grouped in *sori*, on flat or nearly flat receptacles, more rarely scattered over the leaf, typically small, with thin elongated stalks and with small spore output. *Annulus* vertical, incomplete, dehiscence transverse. *Indusium* various or absent.

Family POLYPODIACEAE. *Rhizome* of various habit, typically dictyostelic.

Tribe I. *Woodsieae*. *Sori* globose, often sub-marginal. *Indusium* cup-shaped, opening laterally or splitting irregularly. *Spores* bilateral.

Indusium often ciliate, attached basally. **Woodsia.**

Indusium hooded, attached laterally. **Cystopteris.**

Tribe 2. *Aspidieae*. *Sori* circular or subcircular in outline, flattish; *indusium* attached centrally, peltate.

Sorus cordate, *indusium* reniform, attached in the sinus or absent. **Dryopteris** (including **Phegopteris**).

Sorus circular, *indusium* attached in the centre.

Polystichum.

Tribe 3. *Asplenieae*. *Sori* elongated to linear, developed along veins from which the lateral *indusium* springs.

Subtribe 1. *Blechninae*. *Sori* on veins parallel to the midrib of pinna or pinnule of special fertile leaves.

Only British genus: **Blechnum**.

Subtribe 2. *Aspleninae*. *Sori* on lateral veins of pinna or pinnules of ordinary fronds.

Indusium with single flap entire. **Asplenium**.

Indusium with single flap, margin lacerate.

Athyrium.

Indusium with two flaps, one on either side of sorus.

Scolopendrium.

Tribe 4. *Pterideae*. *Sori* elongated to linear, developed along special veins. *Indusium* mostly absent, sometimes developed as an infolding of the leaf margin.

Subtribe 1. *Gymnogramminae*. *Sori* occupying the whole of the fertile veins.

Only British genus: **Gymnogramme**.

Subtribe 2. *Cheilanthisinae*. *Sori* occupying the distal ends of the fertile veins.

Only British genus: **Cryptogramme**.

Subtribe 3. *Adiantinae*. *Sori* occupying the distal ends of the fertile veins, within the revolute leaf margin.

Only genus: **Adiantum**.

Subtribe 4. *Pteridinae*. *Sori* on intramarginal vein-anastomoses, covered by the indusial leaf margin.

Only British genus (sorus with an additional inner indusial ciliate flap): **Pteridium** (= *Pteris* auct.).

Tribe 5. *Polypodieae*. *Leaves* separating from rhizome by definite absciss-layers. *Indusium* absent.

Only British genus (sori circular): **Polypodium**.

Suborder 2. HYDROPTERIDINEAE. Wall of sporangium without *annulus*. *Spores* of two kinds (heterosporous). *Macrospores* borne singly in macrosporangia, and each developing a female prothallus. *Microspores* numerous in each microsporangium, and developing minute few-celled male prothalli.

Family 1. MARSILIACEAE. *Rhizome of sporophyte* solenostelic with two rows of leaves. *Sori* enclosed in special organs (*sporocarps*) which arise as branches of the leaves. *Female prothallus* producing a single *archegonium*.

Only British genus (leaves filiform): **Pilularia**.

Family 2. SALVINIACEAE. *Rhizome of sporophyte* floating, haplostelic, with two or three rows of partly submerged leaves, on which the *sori*, enclosed in *indusia*, are borne. *Female prothallus* producing several *archegonia*.

Leaves in two rows, deeply bilobed. True roots present†. ***Azolla**.

Order 2. **Ophioglossales**. *Fertile leaves* with one or more simple or pinnate branches bearing the bulky,

† Cf. the exotic genus *Salvinia* which is rootless.

marginal sporangia which arise from groups of cells. *Prothallus* tuberous, subterranean with numerous sunken antheridia and archegonia.

Family OPHIOGLOSSACEAE. *Stem* short, vertical, sunk in the earth and sending up a few leaves which take several years to develop.

Sterile leaf-segment pinnate. Pinnules with digitate nerves. Fertile segments branched. **Botrychium.**

Sterile leaf-segment simple, entire with reticulate veins. Fertile segment unbranched, spike-like.

Ophioglossum.

Division 2. **EMBRYOPHYTA SIPHONOGAMA** (*Phanerogamae auct.*).

Alternation of generations obscured by seed formation. *Sporophyte* cormophytic, very various, heterosporous.

The *megaspore* (*embryo sac*) is not discharged from the *sporangium* (*ovule*) which after fertilisation ripens to form a *seed*.

Microspore (*pollen grain*) on germination forming a *tube*. *Antheridium* represented by a single cell which divides to form two naked *generative cells* which in the lowest Gymnosperms become transformed into *Spermatozoids*.

Subdivision I. **GYMNOSPERMAE.** *Microsporangia* on under surface of sporophylls. *Carpels* not fused together to form an ovary, thus no *stigma* is present.

The *microspores* (*pollen grains*) germinate at the *micropile*. *Seeds* naked on the ovuliferous scales or on a direct prolongation of the axis.

Prothallus (*endosperm*) formed before fertilisation.

Class 1. **CONIFERAE**. *Stem* much branched. No *vessels* in secondary wood. *Leaves* mostly narrow, linear or lanceolate. *Flowers* diclinous. *Perianth* 0 or bracteoid. *Cotyledons* 2-15, free.

Fam. 1. **TAXACEAE**. *Carpels* solitary or few. *Seed* drupaceous exserted.

Subfamily *Taxoideae*. *Stamens* with 3-8 *pollen sacs*. *Carpels* with two ovules or the ♀ *flower* reduced to a single *ovule*.

Tribe *Taxeeae*. ♀ *flower* reduced to a solitary terminal *ovule*.

Only British genus (Mesophyll with no resin canals):
Taxus.

Fam. 2. **PINACEAE**. *Resin canals* always present in leaves. *Micro-* and *megasporophylls* numerous, arranged in cone-like *flowers*. *Seeds* not exserted.

Tribe 1. *Abietineae*. *Leaves* spiral. *Carpels* divided into *cover-scales* and *ovuliferous scales*, the latter with two ovules side by side on the upper surface.

1. Only long shoots present with spiral evergreen leaves.

Bark greyish white. Leaves flat, emarginate with two white lines beneath. Cones erect, scales deciduous.

***Abies.**

Bark reddish. Leaves 4-angled in section, acute. Cones pendulous, scales persistent.

***Picea.**

As *Picea* but leaves flat. ***Tsuga** (including ***Pseudo-tsuga** with long 3-fid cover-scales).

2. Long and short shoots present.

Long and short shoots both with deciduous leaves.
Cone-scales broad, coriaceous, persistent. ***Larix**.

Long and short shoots both with evergreen leaves.
Cone-scales deciduous. ***Cedrus**.

Long shoots with scales, short ones with 2-5 evergreen leaves. Cone-scales woody, persistent. **Pinus**.

Tribe 2. *Cupressineae*. Leaves opposite or whorled.
Ovules erect.

Cone-scales fleshy and confluent in fruit. Only
British genus: **Juniperus**.

Subdivision 2. **ANGIOSPERMAE**. *Ovules* enclosed in an *ovary* formed of coherent *carpels* or of one carpel with coherent margins. *Stigma* present to receive pollen.

Class 1. **MONOCOTYLEDONES**. *Embryo* with one cotyledon. *Mature stem* with closed vascular bundles. *Parallel venation* predominant. *Flowers* often 3-merous.

A. Orders with preponderating inconstancy in the number of floral leaves. (For B see page 23.)

a. Typical achlamydeous (not apopetalous) flowers still occur. (For b see page 21.)

a. Achlamydeous flowers predominate. Great inconstancy in number of stamens and also of carpels. (For β see page 11.)

Order 1. **Pandanales**. Marsh herbs (or tropical trees and lianes). Leaves linear. Inflorescence cylindrical or spherical. Flowers naked or with haplochlamydeous, bracteoid perianth, diclinous, ♂ with $\infty-1$ stamens, ♀ with $\infty-1$ carpels. Seed with endosperm.

Fam. 1. TYPHACEAE. Herbs with rhizomes and distichous, linear leaves. *Inflorescence* cylindrical, ♀ flowers below, ♂ above. ♂ flowers of 2-5 stamens often united; *pollen* in tetrads. ♀ flowers of one carpel on an elongated axis which bears hairs. *Stigma* narrow. *Ovule* one, pendulous. *Fruit* an achene. *Seed* with thin perisperm and fleshy endosperm.

Only genus:

Typha.

Fam. 2. SPARGANIACEAE. Herbs. *Leaves* distichous. *Inflorescence* globose. *Lower inflorescences* ♀, *upper* ♂. *Flowers* homochlamydeous; *tepals* 3-6. *Stamens* 3-6. *Carpels* (1-2) each with one pendulous ovule. *Stigmas* 1-2, lanceolate. *Fruit* a drupe. *Seed* with mealy endosperm.

Only genus:

Sparganium.

β. *Naked flowers still present: but in this order there occur all possible gradations from Achlamydy to Heterochlamydy, also from Hypogyny to Epigyny. Sporophylls definite or indefinite.* (For α see page 10, for γ see page 15.)

Order 2. **Alismatales** (Fluviales, Helobiae). Herbaceous water or marsh plants with *scales* (*Squamulae intravaginales*) in the leaf axils. *Flowers* cyclic or hemicyclic; achlamydeous, haplochlamydeous or diplochlamydeous (homochlamydeous or heterochlamydeous); hypogynous, or epigynous. *Stamens* ∞-1. *Carpels* ∞-1, usually apocarpous when superior. *Endosperm* absent or scanty.

Suborder 1. POTAMOGETONINEAE. *Flowers* hypogynous, achlamydeous or haplo- to homochlamydeous.

Fam. 1. POTAMOGETONACEAE. Submerged or floating fresh or salt water plants. *Leaves* mostly distichous. *Flowers* mostly small, solitary or in spikes, diclinous or monoclinal with 4-1-merous whorls.

Perianth mostly absent. *Stamens* 4-1. *Carpels* 4-1, each with one pendulous ovule. *Fruit* often drupaceous.

a. *Flowers* in spikes.

Tribe 1. *Zostereae*. *Flowers* diclinous in a flat, dorsiventral spike enclosed in the sheath of the uppermost foliage leaf. *Pollen* filiform.

Only British genus :

Zostera.

Tribe 2. *Potamogetoneae*. *Axis of spike* terete, *flowers* monoclinal.

Spike usually many-flowered. *Tepals* 4: *stamens* 4: *achenes* sessile.

Potamogeton.

Spike 1-2-flowered. *Tepals* 0: *stamens* 2: *achenes* stipitate.

Ruppia.

b. *Flowers* solitary or cymose, diclinous.

Tribe 3. *Zannichellieae*. *Perianth* absent or very simple. *Carpels* 4-3.

Only British genus :

Zannichellia.

Fam. 2. NAIADACEAE. Slender, submerged fresh or brackish water herbs. *Stem* with central cylinder of elongated cells enclosing a canal. *Leaves* opposite, linear, toothed, almost imbricating. *Flowers* diclinous. ♂ of two cup-like envelopes and a terminal *anther*. ♀ of one cup-like envelope (which may be absent) and one *carpel* containing one basal anatropous *ovule*.

Only genus :

Naias.

*Fam. 3. APONOGETONACEAE. Herbs with tuberous sympodial stems. *Leaves* submerged or lamina floating. *Spike* aerial enclosed in deciduous *spathe*. *Flowers* monoclinal. *Perianth* 3-1 petaloid. *Stamens* 6 or more. *Carpels* 3-6. *Styles* 3-6. *Fruit* leathery with 2 or ∞ seeds.

Only genus :

***Aponogeton.**

Fam. 4. SCHEUCHZERIACEAE (Juncaginaceae). Scapigerous marsh herbs with narrow, rush-like *leaves*. *Inflorescence* racemose. *Flowers* monoclinal or diclinal, 4-1-merous. *Perianth* homochlamydeous, herbaceous. *Each carpel* with 1-2 anatropous ovules.

Tribe. *Triglochineae*. *All flowers* of inflorescence similar.

Bracts absent. Anthers subsessile, short. **Triglochin**.
 Inflorescence bracteate. Filaments and anthers long.
Scheuchzeria.

Suborder 2. ALISMATINEAE. *Flowers* hypogynous, mostly heterochlamydeous. *Ovules* on the ventral suture.

Fam. 5. ALISMATACEAE. Marsh herbs. *Leaves* chiefly radical. *Inflorescence* often much branched. *Flowers* mostly monoclinal and heterochlamydeous, 3-merous. *Stamens* 6- ∞ . *Carpels* 6- ∞ each with 1- ∞ anatropous ovules. *Styles* 6- ∞ . *Fruit* of achenes or follicles.

a. Flower-axis flat. *Stamens* 6 in one whorl. *Carpels* also more or less cyclic.

a. *Carpels* with one ovule.

Leaves erect, flowers whorled. **Alisma**.

Leaves floating, flowers subsolitary. **Elisma**.

β . *Carpels* with two or more ovules.

Leaves erect, carpels connate. **Damasonium**.

b. Flower-axis convex. *Stamens* mostly more than 6, sometimes spiral. *Carpels* arranged in a head, acyclic.

Flowers hermaphrodite. *Fruit* ribbed, scarcely compressed. **Echinodorus**.

Flowers diclinous. Fruit strongly compressed.

Sagittaria.

Suborder 3. BUTOMINEAE. *Flowers* hypogynous or epigynous, mostly heterochlamydeous. *Ovules* scattered over inner surface of carpel on branching placentae.

Fam. 6. BUTOMACEAE. Marsh plants with conspicuous *flowers*. *Inflorescence* cymose, often umbel-like. *Flowers* monoecious, usually markedly heterochlamydeous, 3-merous. K 3, C 3, A 6 (3 pairs) + 3- ∞ , in the last case the outer ones sterile. G $\overline{6-\infty}$ often united at the base with ∞ ovules scattered over their inner surfaces. *Fruit* of follicles.

Only British genus (perianth homochlamydeous, persistent):

Butomus.

Fam. 7. HYDROCHARITACEAE. Submerged or floating water plants. *Leaves* mostly spiral, sometimes cyclic or distichous. *Flowers* solitary or cymose, at first enclosed in an envelope of 1-2 bracts, usually diclinous, mostly heterochlamydeous, 3-merous. K 3, C 3, A 3 + (3 + 3 + 3 + 3), the inner and outer stamens sometimes staminodal. G $\overline{(2-15)}$. *Placentae* parietal, bearing ∞ orthotropous to anatropous ovules with 2 integuments. *Stigma* often deeply bifid. *Fruit* mostly dehiscent irregularly with ∞ seeds.

a. *Carpels* 6-15. *Placentae* projecting far inwards.

Subfamily 1. *Stratiotoideae*. Fresh water plants. *Foliage leaves* spiral.

Tribe 1. *Stratioteae*. *Leaves* partly submerged. ♀ *flowers* sessile in the 2-leaved spathe.

Leaves erect narrow serrate.

Stratiotes.

Tribe 2. *Hydrochariteae*. Leaves floating. ♀ flowers stalked in the spathe.

Leaves orbicular.

Hydrocharis.

b. Carpels usually 3. *Placentae* not projecting far inwards.

*Subfamily 2. *Vallisnerioideae*. Fresh water plants. *Perianth* often small. *Stigma* short.

*Tribe. *Hydrilleae*. Leaves 1-nerved, in whorls. ♂ flowers 1-3 in a spathe.

Leaves linear, 3 in a whorl.

***Elodea.**

γ. *Naked flowers* dominant. *Stamens* rarely, and *carpels* never, indefinite. (For α see page 10, for β see page 11.)

Order 3. **Glumiflorae**. Grassy herbs with sheathing leaves. Flowers naked, more rarely with perianth of bristles†, in the axils of imbricating bracts (*glumes*). *Gynaecium* always 1-locular with one ovule.

Fam. I. GRAMINACEAE. Herbs (grasses). Stems usually terete. Leaves distichous, sheath usually split to the base and furnished with *ligule* or hairs at junction of sheath and lamina. Flowers in the axils of glumes with 2-nerved 'palea' opposite glume, usually mon-oecious, naked. The lowermost glumes are usually without flowers (*empty*). Deeply bifid scale (*lodicaulae*) often present opposite palea. Stamens mostly 3. Anthers versatile. Stigmas usually 2, feathery. Fruit a caryopsis‡. Seed with copious endosperm. Embryo with peltate enlargement (*scutellum*) of cotyledon in whose distal

† *Oreobolus*, a South American genus of *Cyperaceae*, tribe *Rhynchosporae*, has a perianth of glume-like tepals.

‡ A caryopsis is a superior achene in which the pericarp and testa are united.

hollow lie the plumule and the radicle, the latter surrounded by a sheath (the *Coleorrhiza*).

Subfamily 1. *Panicoideae*. *Spikelets* 1-flowered with axis (*rachilla*) not produced beyond flower, rarely 2-flowered, in which case the lower flower is imperfect, when ripe falling off as a whole. (See footnote.)

a. Hilum point-like, *spikelets* dorsally compressed or terete.

Tribe 1. *Paniceae*. *Flowering glumes* and *paleae* harder than *empty glumes*. *Lower empty glume* usually smaller.

Pedicels of spikelets naked or hairy. ☉ *Panicum*.

Pedicels of spikelets with stiff bristles. ☉ *Setaria*.

b. Hilum linear, *spikelets* laterally compressed.

Tribe 2. *Oryzeae*. *Empty glumes* minute or 0. *Stamens* often 6 (but 3 in Brit. Sp.).

Empty glumes absent. Only British genus: *Leersia*.

Subfamily 2. *Poëoideae*. *Spikelets* 1-many-flowered, when 1-flowered the axis (*rachilla*) is often prolonged beyond the flower, mostly jointed distally to the *empty glumes* which remain behind when the *flowering glumes* fall away†. When 2-many-flowered there are always distinct internodes between the flowers.

a. Spikelets with distinct stalks, arranged in panicles, spike-like panicles or racemes. (For β see page 19.)

Tribe 3. *Phalarideae*. *Spikelets* 1-flowered, with 4 *empty glumes*. *Palea* 1-nerved.

3rd and 4th glumes empty, awnless, reduced to small scales. *Phalaris*.

† In *Alopecurus*, *Polypogon*, *Holcus* and *Spartina* the spikelets fall as a whole.

3rd and 4th glumes empty, small with dorsal awns.

Anthoxanthum.

3rd and 4th glumes, or at any rate the 3rd with a triandrous ♂ flower, almost as long as the 1st and 2nd. Upper flowers hermaphrodite, diandrous. **Hierochloë.**

Tribe 4. *Agrostideae*. *Spikelets* 1-flowered with two *empty glumes*. *Palea* 2-nerved.

* Rachilla not produced beyond flowering glume.

Panicle effuse, fruit included in hardened glume.

Milium.

Panicle dense, cylindric, spikelets falling as a whole, flowering glume with dorsal, bent awn. **Alopecurus.**

Panicle dense, cylindric, empty glumes persistent, flowering glume awnless. **Phleum.**

Spikelets dorsally compressed in a simple spike.

Mibora.

Panicle loose, flowering glume small, membranous.

Agrostis.

Panicle contracted, spikelets falling as a whole, empty glumes with awns. **Polypogon.**

Rachilla with long silky hairs, empty glumes acuminate. **Calamagrostis.**

** Rachilla produced beyond the flowering glume.

Empty glumes large, dilated at base, much larger than the minute 4-toothed flowering glume. **Gastridium.**

Empty glumes large, flowering glumes bifid, with slender awns. **Apera.**

Empty glumes large, flowering glumes with awns, rachilla ciliate. **Deyeuxia.**

Spikelets large in dense panicle, rachilla long and silky. **Ammophila.**

Empty glumes with feathery hairs, flowering glume 3-awned. **Lagurus.**

Tribe 5. *Aveneae*. *Spikelets* 2-many-flowered. *Flowering glumes* usually shorter than the two *empty glumes*, usually with dorsal, rarely apical, bent and twisted awn.

A. Spikelets falling as a whole. **Holcus.**

B. Flowering glumes in fruit becoming detached from the persistent empty glumes.

α. Rachilla not produced.

Flowering glumes awned, 2-toothed. **Aira.**

β. Rachilla produced.

Flowers 2, awn bent in middle, tip clavate.

Corynephorus.

Flowers 2, awn straight, acute. **Deschampsia.**

Flowers 2-6, flowering glumes deeply bifid with twisted awns. **Trisetum.**

Flowers 2-6, flowering glumes entire or 2-toothed with long awns. **Avena.**

Flowers 2, upper hermaphrodite, lower ♂.

Arrhenatherum.

Tribe 6. *Festuceae*. As *Aveneae* but *flowering glumes* usually longer than the two *empty glumes*. *Awn* absent, or apical and straight.

α. Spikelets 2 or more flowered: rachilla not bearded. Flowering glumes with three broad teeth.

Sieglingia.

β. Spikelets 2 or more flowered: rachilla with silky hairs. **Phragmites.**

γ. Spikelets subspicate or capitate, with imperfect spikelets on pedicels below them.

Imperfect spikelets ciliate.

Sesleria.

Imperfect spikelets with stiff bristles.

Cynosurus.

δ. Spikelets 2 or more flowered. Flowering glumes 1 or 3-nerved, all alike.

Spikelets arranged in a spike-like panicle. Flowering glumes scarious. **Koeleria.**

Spikelets conical, terete, in a slender panicle. Flowering glumes cartilaginous. **Molinia.**

Spikelets in effuse panicle, branches whorled. Flowering glumes coriaceous. **Catabrosa.**

ε. Spikelets 2 or more flowered. Flowering glumes 3-5-nerved, upper empty, convolute. **Melica.**

ζ. Spikelets 3-many-flowered. Flowering glumes 5 or more nerved.

Spikelets few-flowered, clustered in secund panicles. **Dactylis.**

Spikelets paniced, drooping. Glumes broad, obtuse. **Briza.**

Spikelets paniced. Flowering glumes compressed, keeled, tips nerved, without awns. **Poa.**

Spikelets very many flowered. Flowering glumes convex, obtuse, tips without nerves or awn. **Glyceria.**

Spikelets in racemes or spike-like panicles. Flowering glumes convex, tip nerved, acute or with awn. Ovary glabrous. **Festuca.**

Spikelets many-flowered, in panicles. Flowering glumes convex. Ovary tip villous. **Bromus.**

Spikelets sessile, distichous. Flowering glumes convex. Ovary tip villous. **Brachypodium.**

β. *Spikelets* in two rows which approach one another forming a unilateral spike or raceme with unjointed axis. (For α see page 16.)

Tribe 7. *Chlorideae*. *Spikelets* 1-many-flowered. *Empty glumes* and *paleae* mostly 2-nerved.

Spikelets falling as a whole. Spikes solitary.

Spartina.

Empty glumes persistent. Spikes digitate. **Cynodon.**

γ . *Spikelets* in two opposite rows. (For α see page 16, for β see page 19.)

Tribe 8. *Hordeae*. *Spikelets* sessile in notches on a simple *rachis* forming a *spike*.

α . Style 1. Spikes unilateral.

Nardus.

β . Styles 2. Spikes bilateral.

I. Spikelets solitary in notches of rachis.

† Spikelets with the narrow sides to rachis. **Lolium.**

†† Spikelets inserted broadside to the rachis.

Spikelets 1-2-flowered in a slender spike. Spikelets sunken in the notches. **Lepturus.**

Spike stout. Spikelets not sunken in the notches.

Triticum (incl. **Agropyrum** and ***Secale**).

II. 2-6 spikelets in each notch.

1. Spikelets 1-flowered.

Hordeum.

2. Spikelets 2-many-flowered.

Elymus.

Fam. 2. CYPERACEAE. Grass-like herbs often with triangular stems. *Leaves* with closed sheaths. *Ligule* usually absent. *Flowers* in spikelets or spikelet-like cymes, monoclinal or diclinal, naked or rarely with homochlamydeous perianth. *Stamens* mostly 3-1, rarely more. *Anthers* basifixed. *Carpels* (3-2). *Styles* 3-2 with filiform *stigmas*. *Fruit* an achene with free seed.

Subfamily 1. *Scirpoideae*. *Spikelets* with many hermaphrodite flowers. Individual flowers in the spikelet may be diclinal. *Perianth* present or absent.

Tribe 1. *Scirpeae*. No *bracteoles*.

Subtribe 1. *Cyperinae*. *Glumes* distichous.

Only British genus :

Cyperus.

Subtribe 2. *Scirpinae*. *Glumes* spiral.

Spikelets solitary, terminal. Bristles 3-8, included.

Style thickened at base.

Heleocharis.

Spikelets clustered and lateral. Bristles 0, or 3-8, included. Style not thickened at base.

Scirpus.

Spikelets solitary or clustered, terminal. Bristles very long, cottony.

Eriophorum.

Subfamily 2. *Rhynchosporoideae*. *Flowers* in spikelet-like cymes (*pseudo-spikelets*) which are arranged in spikes or heads, mono- or diclinous. *Perianth bristles* present or absent.

Tribe 2. *Rhynchosporeae*. *Pseudo-spikelets* few-flowered.

Pseudo-spikelets terete. Bristles slender or absent.

Fruit beaked.

Rhynchospora.

Pseudo-spikelets compressed; glumes distichous.

Fruit not beaked.

Schoenus.

Pseudo-spikelets terete. Bristles 0. Fruit obtuse.

Cladium.

Subfamily 3. *Caricoideae*. *Flowers* always naked, diclinous, in many-flowered (rarely few-flowered), mono- or diclinous spikelets. ♀ *flower* enclosed in modified bracteole (*utriculus*).

Spikelets 1-2-flowered.

Kobresia.

Spikelets many-flowered.

Carex.

b. Achlamydeous flowers rare: their occurrence is mostly due to reduction and is correlated with spathe-development. Definite number of stamens and carpels

dominant, but numerous stamens and more than three carpels often occur. (For a see page 10.)

Order 4. **Spathiflorae.** Mostly sympodial, rarely forming erect stems. (The *Lemnaceae* are free floating and undifferentiated.) *Flowers* cyclic, haplochlamydeous or diplochlamydeous, homochlamydeous or naked, 3-2-merous, hermaphrodite or diclinous, often very reduced, always inserted on a *spadix* more or less enveloped by a bract (*spathe*). *Floral bracts* absent.

Fam. 1. ARACEAE. Mostly herbs. *Rhizome* often tuberous. *Leaves* often with netted veins. *Flowers* mono- or diclinous, rarely dioecious, 2-3-merous, or reduced sometimes to a single *stamen* or *carpel*. *Fruit* usually a berry. *Seed* with fleshy outer integument.

Subfamily 1. *Pothoideae*. Land plants. No *latex* or *raphides* present. *Leaves* distichous or spiral. Lateral nerves of 2nd and 3rd order usually netted. *Flowers* mostly hermaphrodite.

Tribe 1. *Acoreae*. *Leaves* not differentiated into stalk and lamina. *Perianth* present. *Ovules* orthotropous.

Only British genus :

☉ **Acorus.**

Subfamily 2. *Aroideae*. Mostly tuberous land or marsh plants with netted leaves. *Latex* present. *Flowers* diclinous. *Perianth* usually absent.

Tribe 2. *Areae*. *Spadix* mostly with vestigial flowers. *Gynaecium* of one carpel.

Only British genus :

Arum.

Fam. 2. LEMNACEAE. Small free-swimming thalloid plants not differentiated into *stem* and *leaves*. *Flowers* monoecious. ♂ flower of one *stamen*. ♀ of one *carpel* with 1-6 basal *ovules*.

Subfamily 1. *Lemnoideae*. *Roots* present. *Inflor-escence* with spathe and two ♂ flowers.

Roots several.

Spirodela.

Root solitary.

Lemna.

Subfamily 2. *Wolffoideae*. *Roots* absent. *Inflor-escence* without spathe and with only one ♂ flower.

Wolffia.

B. Orders with typically pentacyclic flowers. Whorls typically isomeric, mostly 3-merous.
(For A see page 10.)

a. Flowers homochlamydeous to heterochlamydeous, very rarely naked. Bracteoid perianths still occur but petaloid perianths predominate. Hypogyny and Actinomorphy preponderate. (For *b* see page 29.)

Order 5. **Farinosae**. Mostly herbs, rarely with well-developed stem. *Flowers* cyclic, homo- or heterochlamydeous, 3 or 2-merous. General floral formula $T_3 + T_3$ (more rarely $K_3 + C_3$), $A_3 + 3$, $G(3)$. One whorl, or all but one of the stamens may be absent. *Ovules* orthotropous, but anatropous ovules also occur. *Endosperm* mealy.

Fam. ERIOCAULACEAE. Perennial, scapigerous herbs. *Leaves* narrow, chiefly radical. *Flowers* in involucrate heads, monoclinal, usually monoecious. *Perianth* membranous. Outer whorl of stamens mostly absent. *Carpels* (2-3). *Styles* 2-3. *Ovary* 2-3-loc, each locus

with a pendulous orthotropous ovule. *Fruit* a loculicidal capsule.

Subfamily *Eriocauloideae*. *Stamens* 4 or 6. *Perianth-segments* free, inner ones with gland near apex.

Only British genus :

Eriocaulon.

Order 6. **Liliiflorae.** Characters of *Farinosae*, but *ovules* mostly anatropous, and *seeds* usually with fleshy or cartilaginous endosperm. Exceptionally 2 or 4-merous flowers occur.

Suborder 1. JUNCINEAE. *Perianth* homochlamydeous, not petaloid. *Endosperm* containing starch.

Fam. 1. JUNCACEAE. Herbs, usually perennating by underground sympodial rhizomes. *Inflorescence* compound, usually many-flowered. *Flowers* homochlamydeous, 3-merous, mostly monoclinal. *Perianth* bract-like. The inner whorl of *stamens* sometimes not developed. *Carpels* (3). *Style* 1 with three filiform *stigmas*. *Ovary* 1 or 3-loc with 1 or ∞ ovules. *Capsule* loculicidal. *Embryo* straight.

Annual or perennial. Glabrous. Capsules 3-loc, many-seeded. **Juncus.**

Perennial. Hairy. Capsule 1-loc, 3-seeded. **Luzula.**

Suborder 2. LILIINEAE. *Perianth* mostly petaloid, homo- very rarely heterochlamydeous. Inner circle of *stamens* present. *Endosperm* with no starch.

Fam. 2. LILIACEAE. *Flowers* usually homochlamydeous and monoclinal. *Perianth* usually petaloid, *segments* free or united. *Stamens* 6 (3 in *Ruscus*). *Ovary* usually superior, mostly 3-loc with axile *placentae*. *Fruit* various.

Subfamily 1. *Melanthioideae*. *Rhizome* or *corm*.
Inflorescence terminal. *Anthers* extrorse and *capsule* septicidal, or *anthers* introrse and *capsule* septicidal, or *anthers* extrorse and *capsule* loculicidal, rarely *anthers* introrse and *capsule* loculicidal. *Fruit* never a berry.

a. *Rhizome*. *Seeds* long, flat and winged, or angled.

Tribe 1. *Tofieldieae*. *Leaves* sessile, distichous.
Stamens 6. *Anthers* introrse. *Style* divided or 0.

Flowers yellow. *Style* very short, stigma small.
Capsule loculicidal. **Narthecium.**

Flowers greenish-white. *Style* absent, stigmas 3, short. *Capsule* septicidal. **Tofieldia.**

b. *Corm*, or short *rhizome*. *Seeds* subglobose.

Tribe 2. *Colchiceae*. *Leaves* radical. *Scape* short subterranean with 1-3 leaves. *Anthers* introrse. *Capsule* septicidal.

Perianth tube long. *Styles* separate. **Colchicum.**

Subfamily 2. *Asphodeloideae*. *Rhizome*. *Leaves* usually radical. *Inflorescence* mostly terminal. *Anthers* introrse, sometimes opening at apex. *Capsule* loculicidal.

Only British genus : **Simethis.**

Subfamily 3. *Allioideae*. *Bulb* or short *rhizome*.
Inflorescence a cymose umbel enclosed by two broad, sometimes united, leaves, more rarely subtended by two narrow bracts, or reduced to a single flower.

Tribe 3. *Allieae*. *Bulb*, or stem with thickened base. *Perianth* free or united. *Stamens* 6 or only 3 fertile.

Inflorescence few-flowered, subtended by two free bracts. **Gagea.**

Inflorescence usually many-flowered, subtended by two coriaceous, mostly united, bracts. **Allium.**

Subfamily 4. *Lilioideae*. *Bulb*. *Inflorescence* terminal, racemose. *Anthers* introrse. *Capsule* loculicidal.

Tribe 4. *Tulipeae*. *Bulb*, imbricated or tunicated. *Stem* bearing several foliage leaves, rarely only one. *Flowers* few, axillary, or solitary and terminal.

a. *Anthers* versatile.

Flowers large. *Nectary* median or obscure. ***Lilium.**

β. *Anthers* basifixed.

Flowers large, drooping, segments not reflexed. *Nectary* oblong. **Fritillaria.**

As *Fritillaria* but flowers erect and nectary absent.

Tulipa.

Flowers few, small. *Perianth-segments* spreading.

Lloydia.

Tribe 5. *Scilleae*. *Tunicated bulb*. *Aerial stem* without leaves. *Flowers* in the axils of bracts.

1. *Perianth-segments* free or nearly so.

Filament filiform or flattened at base. **Scilla.**

Filaments flattened throughout. **Ornithogalum.**

2. *Perianth-segments* united.

[*Corolla* funnel-shaped, not contracted, all flowers fertile. ***Hyacinthus.**]

Corolla globose, upper flowers imperfect. **Muscari.**

Subfamily 5. *Asparagoideae*. *Rhizome* subterranean, ending in flowering shoots, or monopodial with lateral flowering shoots. *Fruit* a berry.

a. *Flowers* homochlamydeous.

Tribe 6. *Asparageae*. *Rhizome* ending in leafy stems. *Cauline leaves* small, scale-like, subtending narrow or broad leaf-like shoots (*Phylloclades*).

Stem herbaceous. Flowers axillary. Stamens 6. Filaments distinct. **Asparagus.**

Stem shrubby. Flowers on the phylloclades dioecious. Stamens 3 with connate filaments. **Ruscus.**

Tribe 7. *Polygonateae*. As *Asparageae*, but *stem* with large, broad *foliage leaves*.

Leaves many. Flowers axillary. Perianth tubular 6-cleft. **Polygonatum.**

Leaves 2. Flowers in terminal racemes. Perianth 4-partite. **Maianthemum.**

Tribe 8. *Convallarieae*. *Rhizome* monopodial with lateral *inflorescence-axes*.

Subtribe *Convallariinae*. Style ending in one small stigma. **Convallaria.**

b. Flowers heterochlamydeous.

Tribe 9. *Parideae*. *Foliage leaves* net-veined forming a whorl. *Flowers* solitary or umbellate. *Fruit* a berry.

Leaves 4 or more in a whorl. Flowers 4 or more merous. Only British genus: **Paris.**

Fam. 3. AMARYLLIDACEAE. Resembling, in essential points, Liliaceae. *Anthers* mostly introrse. A *corona*, formed of staminal stipules, is often present. *Ovary* mostly inferior. *Placentation* axile. *Ovules* anatropous in two rows on each placenta. *Fruit* a loculicidal capsule or berry. *Seeds* usually few.

Subfamily 1. *Amaryllidoideae*. *Bulb*. *Scape* leafless with a solitary flower, or involucrate umbellate inflorescence.

Tribe 1. *Amaryllideae*. No *corona*.

Subtribe *Galanthinae*. Perianth actinomorphic, without tube. Loculi of ovary with ∞ ovules.

Perianth-segments all equal.

Leucojum.

Outer perianth-segments larger than inner.

☉ **Galanthus.**

Tribe 2. *Narcisseae*. *Corona* present as tube, ring or scales.

Subtribe *Narcissinae*. Stamens included in the cup-like *corona*.

Narcissus.

Fam. 4. DIOSCOREACEAE. Lianes. *Rhizome* often tuberous. *Leaves* often sagittate, veins reticulate. *Flowers* in racemes, mostly diclinous, homochlamydeous. *Perianth* sepaloid, usually with short tube. $G(\bar{3})$, 3 or 1-loc with axile or parietal *placentae*, each placenta mostly with two superposed, anatropous ovules. *Styles* 3, sometimes bifid. *Fruit* a capsule or berry.

Tribe *Dioscoreae*. *Flowers* diclinous. *Ovules* 2 in each loculus.

Only British genus (berry imperfectly 3-celled):

Tamus.

Suborder 3. IRIDINEAE. As Liliineae, but *inner whorl of stamens* aborted.

Fam. 5. IRIDACEAE. Mostly herbs with equitant leaves and terminal inflorescences. *Flowers* hermaphrodite, homo- or heterochlamydeous 3-merous, actinomorphic or zygomorphic. *Stamens* always 3 (the outer whorl). *Anthers* extrorse. $G(\bar{3})$ with many anatropous ovules

on axile placentae†. *Styles* 3, often divided and leaf-like. *Capsule* loculicidal with globose or angular seeds.

Subfamily 1. *Crocoideae*. Small herbs often with underground flower stalks. *Flowers* solitary, or several axillary around a terminal flower.

Scape short. Perianth tube short. **Romulea.**

Scape 0. Perianth tube long. **⊙Crocus.**

Subfamily 2. *Iridoideae*. *Stem* distinct. *Leaves* equitant. *Flowers* several in each spathe.

Tribe 1. *Moraeae*. *Perianth tube* short or absent. *Stigmas* on underside of leaf-like style-arms. *Capsule* not enclosed in spathe.

Only British genus: **Iris.**

Tribe 2. *Sisyrinchieae*. As above, but *style-arms* mostly terete.

Only British genus: **⊙Sisyrinchium.**

Subfamily 3. *Ixioideae*. As *Iridoideae*, but only *one flower* enclosed in each *spathe*.

Tribe *Gladioleae*. *Flowers* strongly zygomorphic, often curved.

Only British genus: **Gladiolus.**

b. Flowers homochlamydeous to heterochlamydeous but in the first case the perianth is petaloid. Epigyny throughout. Zygomorphy dominant. (For a see page 23.)

Order 7. **Microspermae**. *Flowers* cyclic, homo- or heterochlamydeous, 3-merous, typically diplostemonous but *androecium* often shows marked reduction.

† The Mediterranean genus *Hermodactylus* has a 1-loc ovary with parietal placentae. In all other respects this monotypic genus is very near to *Iris*.

Ovary inferior, 3 or 1-loc with numerous, minute ovules. *Endosperm* present or absent.

Suborder GYNANDRAE. *Flowers* always zygomorphic, no *endosperm*.

Family ORCHIDACEAE. Perennial herbs. *Flowers* homo- or heterochlamydeous, typically 3-merous, almost invariably monoclinal, zygomorphic, usually resupinate by twisting of the ovary so that the posterior petal occupies an anterior position and is often developed into a labellum. Of the *stamens* usually only the anterior† one of the outer whorl is fertile. In subfamily Diandrae the two lateral stamens of the inner whorl are usually fertile. Frequently (e.g. in *Orchis*) the two anterior stamens of the inner whorl are represented by staminodes on the sides of the column. *Pollen* cohering in masses (*Pollinia*). *Carpels* ($\bar{3}$) sunk in the floral axis which is prolonged beyond the insertion of the perianth into a *column* which bears the stamens. *Stigmas* 3, usually on the surface of the column, the posterior one rudimentary or developed into a beak-like structure (*rostellum*) beneath the anther or between its cells. *Ovary* mostly 1-loc with 3 parietal placentae and ∞ ovules. *Fruit* a capsule with very numerous, minute seeds without *endosperm*. *Embryo* not, or only slightly, differentiated.

Subfamily 1. *Pleonandrae* (Diandrae). The two lateral (rarely all 3) stamens of the inner whorl fertile. All three stigmas similar and receptive.

† The terms *anterior* and *posterior* in this description refer to position before the occurrence of resupination.

Tribe 1. *Cypripedileae*. Flowers zygomorphic. Column curved towards the slipper-shaped labellum.

Only British genus: **Cypripedium**.

Subfamily 2. *Monandreae*. Anterior stamen of outer whorl fertile. Lateral stigmas receptive, odd one rudimentary or developed as rostellum.

a. *Basitonae*. Anther confluent with column. Pollinia with appendages at base which are in connection with glands in rostellum.

Tribe 2. *Ophrydeae*. The only tribe. Terrestrial orchids with root tubers.

Subtribe *Serapiadinae*. Column short with labellum at base. Anthers erect. Stigma disc-like. Glands of pollinia enclosed in rostellar pouch.

Spur long: both glands in a single pouch. **Orchis**.

Spur absent: both glands in one pouch. **Aceras**.

Spur absent: glands in two distinct pouches.

Ophrys.

Subtribe *Gymnadeniinae*. As above, but pollen masses naked or surrounded by processes of the anthers.

Without spur. **Herminium**.

With spur. **Habenaria**.

b. *Acrotonae*. Anther deciduous. Pollen masses without appendages.

(α) *Acranthae*: Inflorescences terminal on branches whose basal portions make up a sympodial shoot.

[(β) *Pleuranthae* (main axis sympodial but flowers borne on special lateral axes): not represented in Britain.]

I. *Convolutae*. Leaves convolute in bud. Sheath and blade not distinct from one another. Pollen granular. Anther withering without falling.

Tribe 3. *Neottieae* (only tribe).

Subtribe 1. *Cephalantherinae*. *Labellum* with distinct proximal lobe (*hypochilium*) which is often spurred. *Anthers* erect. *Rostellum* short or absent.

α. Stem leafy: no spur.

Flowers in racemes, ovary straight. **Epipactis.**

Flowers in spikes, ovary twisted. **Cephalanthera.**

β. Stem scaly, without foliage leaves, *labellum* spurred.

Labellum turned upwards, column short. **Epipogon.**

Subtribe 2. *Spiranthinae*. *Leaves* soft, net-veined. *Anther* as long as the beaked *rostellum* and lying against it. *Pollinia* not divided into distinct masses.

A. Median sepal and petals forming a helmet surrounding the base of the *labellum*.

Spike spirally twisted. **Spiranthes.**

B. Sepals and petals standing apart.

Scape with two opposite leaves. **Listera.**

Scape brown, leafless. **Neottia.**

Subtribe 3. *Physurinae*. As above but *pollinia* divided into numerous distinct masses.

Rhizome creeping, leaves several. **Goodyera.**

II. *Duplicatae*. *Leaves* folded in bud.

Tribe 4. *Liparideae*. *Pollinia* 4, waxy, without appendages.

A. Foliage leaves present.

Column short, straight. Anther persistent. **Malaxis.**

Column slender, bent forwards. Anther deciduous.

Liparis.

B. No foliage leaves. Roots absent. Rhizome branched, coral-like. **Corallorhiza.**

Class 2. **DICOTYLEDONES.** *Embryo* usually with two cotyledons. *Stem* with open bundles. *Leaves* usually with reticulate venation. *Flowers* often 4-5-merous.

Subclass 1. **ARCHICHLAMYDEAE.** *Perianth* simple or absent. *Flowers* either achlamydeous, or haplochlamydeous, or diplochlamydeous with polypetalous corolla. Sympetaly rare (marked in *Cotyledon*). Apopetaly occurs not infrequently.

a. **Amentiflorae** (Moss)†. **Mostly trees or shrubs.** ♂ flowers usually, and ♀ flowers often, in catkins. *Perianth* absent or haplochlamydeous and bracteoid. (For *b* see p. 37.)

Order 1. **Salicales.** Deciduous trees or shrubs. *Leaves* alternate, stipulate, seldom deeply lobed. *Flowers* dioecious, both sexes in simple catkins. *Perianth* cup- or saucer-shaped, or modified into one or two or rarely more nectaries. *Stamens* 2-∞, filaments free or rarely connate. *Carpels* (2). *Ovary* 1-locular with ∞ ovules on two parietal placentae. *Ovules* anatropous with two integuments. *Fruit* a loculicidal capsule quite free from the bracts. *Seeds* small with basal tuft of hairs. *Endosperm* 0.

except 5
purple
or casto

Only family: **SALICACEAE.**

† Engler has no names for these groups, simply referring to them as *a*, *b*, *c* and *d*. The names used in this book are those made use of by Dr Moss in his lectures at Cambridge.

Leaves usually broad. ^{long petiole} Bracts laciniate. [^] Perianth cup-like or saucer-like, more or less oblique. **Populus.**

Leaves usually narrow. ^{short petiole} Bracts entire. Perianth of usually separate nectaries. [^] Catkins erect. **Salix.**

Order 2. **Myricales.** Shrubs or trees. Leaves alternate, simple, exstipulate, sweet scented. *Flowers* achlamydeous, diclinous, monoecious or dioecious, sometimes with bracteoles at the base. *Stamens* usually 4. *Carpels* (2). *Ovary* 1-locular with one basal orthotropous ovule with one integument. *Styles* 2, filiform. *Fruit* a drupe with waxy surface often adnate to bracteoles. *Seeds* without endosperm.

Fam. MYRICACEAE.

Only British genus: (no connection with real ^{Myrica} **Myrica** ^{gale} _(Bog myrtle))

Order 3. ***Juglandales.** Trees with alternate, exstipulate, pinnate leaves. *Flowers* diclinous, monoecious, achlamydeous or haplochlamydeous. *Stamens* 3-40. *Carpels* (2). *Ovary* inferior, 1-locular with one basal, orthotropous ovule with one integument. *Fruit* a drupe or nut with adherent bracts. *Endosperm* 0.

Fam. ***JUGLANDACEAE.** ♀ *flower* with perianth which is fused with the ovary and also with the bract and bracteoles. (= wing).

Walnut = *J. regia* (Nat of E. Europe)

***Juglans.**

Order 4. **Fagales.** Trees or shrubs. *Leaves* alternate, simple, with stipular, usually caducous bud-scales. *Flowers* homochlamydeous or achlamydeous, mostly diclinous and monoecious. ♂ *flowers* in catkins. ♀ *flowers* in catkins or small spikes, or 1-3 seated in an involucre of free or connate bracts. *Stamens* often

opposite the tepals. *Carpels* (2-6). *Ovary* sub-inferior, usually more or less completely 2-3-locular (after fertilisation) with 1-2 pendulous, anatropous, parietal ovules in each loculus. *Fruit* usually a nut. *Endosperm* 0.

Fam. 1. BETULACEAE. Flowers usually appearing before leaves. *Perianth* present in flowers of only one sex. ♂ *catkins* compound. ♂ *flowers* united to their bracts. *Stamens* 2-10. *Filaments* often dividing into two, each branch bearing a half-anther. ♀ *catkins* often minute. *Carpels* (2). *Styles* 2.

Tribe 1. *Coryleae*. ♂ *flowers* without perianth. ♀ *flowers* with perianth. *Fruit* not winged, enclosed in the enlarged, herbaceous bracts.

Leaves in bud folded parallel to lateral nerves. ♀ spike large. Fruiting bracts open 3-lobed. **Carpinus.**

Leaves in bud folded parallel to midrib. ♀ spike minute. Fruiting bracts forming a cupule. **Corylus.**

Tribe 2. *Betuleae*. ♂ *flowers* in dichasia, with perianth. ♀ *flowers* without perianth. *Fruit* usually winged.

Filaments branched. ♀ *catkins* falling at end of the first summer. **Betula.**

Filaments simple. Empty ♀ *catkins* persistent for more than one year. **Alnus.**

Fam. 2. FAGACEAE. *Flowers* and *leaves* appearing together, or flowers appearing after leaves. *Flowers* of both sexes with perianth of 4-7 connate tepals. ♂ *catkins* simple or compound. *Stamens* 4-14. ♀ *flowers* in 3-2-flowered dichasia or solitary. *Carpels* usually 3-4. *Styles* 3-4. *Fruit* surrounded by involucre (*cupule*).

Tribe 1. *Fageae*. ♂ *catkins* globose. Two angular *nuts* in each cupule.

Only British genus :

Fagus.

Tribe 2. *Castaneae*. ♂ *catkins* elongate. *Nuts* rounded.

♂ *catkins* erect. *Nuts* 3, enclosed in spinous cupule.

⊙ **Castanea.**

♂ *catkins* pendulous. Nut single, exserted. **Quercus.**

Order 5. **Urticales.** Herbs, shrubs or trees with stipulate, often roughly hairy, leaves. *Inflorescence* mostly cymose, but *catkins* still occur (*Urtica*). *Flowers* cyclic, homochlamydeous, rarely naked. *Stamens* opposite the tepals. *Carpels* 2-1, superior. *Ovary* usually 1-locular, with one ovule with two integuments.

Fam. 1. **ULMACEAE.** Woody plants without latex. *Leaves* often distichous, simple, oblique at base, strongly pinnately nerved, serrate, with rough hairs and deciduous stipules. *Flowers* mostly hermaphrodite. *Perianth* 4-5-lobed. *Stamens* as many. *Filaments* straight in bud. *Carpels* (2). *Styles* 2. *Ovule* usually solitary, pendulous and anatropous. *Fruit* a nut or drupe. *Endosperm* usually 0.

Subfamily *Ulmoideae*. *Flowers* in clusters. *Peduncles* in the axils of scales. *Fruit* never drupaceous. *Embryo* straight.

Only British genus (fruit broadly winged): **Ulmus.**

Fam. 2. **CANNABACEAE.** Aromatic herbs without latex. *Leaves* palmately nerved, and lobed or divided. *Stipules* persistent. *Flowers* dioecious. ♂ *flowers*, *tepals* 5, *stamens* 5. ♀ *flowers* with perianth of low entire mar-

gined cup surrounding the *gynaecium*. *Carpels* 2. *Styles* 2. *Ovule* pendulous. *Fruit* a nut. *Endosperm* fleshy. *Embryo* curved.

Stem twining. Leaves cordate, 3-7-lobed. **Humulus.** *Hop*

Stem erect. Leaves palmate. ***Cannabis.** *Hemp*

Fam. 3. URTICACEAE. Mostly herbs without latex. *Stinging hairs* often present. *Flowers* diclinous. *Perianth* of both sexes usually of four free tepals. *Stamens* as many incurved in bud. *Carpel* 1. *Fruit* a nut enclosed by inner two or by all the tepals. *Endosperm* o. *Embryo* straight.

Tribe 1. *Urereae*. *Stinging hairs* present. *Perianth* of ♀ flowers 4-partite.

Only British genus (leaves opposite, stigma brush-like): **Urtica.**

Tribe 2. *Parietarieae*. No *stinging hairs*. ♀ *perianth* (or involucre?) tubular.

Only British genus: **Parietaria.**

b. **Petaloidae** (Moss). **Flowers** haplochlamydeous, **perianth** often petaloid. **Diplochlamydeous flowers** rare (**Rumex**). (For *a* see page 33, for *c* see page 40.)

Order 6. **Santalales**. Partial or entire parasites. *Leaves* entire, exstipulate. *Flowers* cyclic, homochlamydeous with *stamens* opposite *tepals*. *Carpels* (2-3), rarely (1). One *ovule* to each carpel, pendulous from apex of loculus or from free central placenta, or placenta and ovules not differentiated. One or no integuments.

Suborder I. SANTALINEAE. *Ovules* differentiated from placenta, often without integument.

Fam. SANTALACEAE. Chlorophyllous hemi-parasites. *Leaves* mostly alternate, entire, exstipulate. *Tepals* 4-5, valvate in bud, bracteoid or petaloid, united below. *Stamens* adnate to tepals. *Ovary* 1-locular with a columnar, free central placenta from which are suspended the 1-3 ovules which are without integument. *Fruit* indehiscent, with one endospermous seed.

Tribe *Thesieae*. *Perianth* epigynous. *Tube* long.

Only British genus :

Thesium.

Suborder 2. LORANTHINEAE. *Ovules* mostly not differentiated.

Fam. LORANTHACEAE. Evergreen shrubs, mostly parasitic on trees. *Leaves* thick, usually opposite and exstipulate. *Tepals* usually 4, valvate, fleshy. *Floral axis* united to *gynaecium*, often forming a crenate ring (*calyculus*). *Ovary* 1-loc, usually with only one fertile embryo sac. *Fruit* a pseudo-berry, inner layer of axis becoming sticky.

Subfamily *Viscoideae*. *Flowers* diclinous. No distinct *calyculus*.

Tribe *Visceae*. *Flowers* single or in groups in the axils of persistent bracts. *Placenta* basal.

Only British genus :

Viscum. (*Mistle*)

Order 7. **Aristolochiales**. *Flowers* cyclic, homochlamydeous, epigynous, actinomorphic or zygomorphic. *Perianth* petaloid. *Ovary* inferior, usually 3-6-locular. *Ovules* always ∞ , mostly axile.

Fam. ARISTOLOCHIACEAE. Herbs or lianes, with alternate, long-stalked, usually cordate or reniform, exstipulate leaves. *Flowers* usually hermaphrodite, actinomorphic or zygomorphic. *Perianth* petaloid, with three

lobes or one lip. *Stamens* usually 6 or 12, free or united to the style. *Carpels* ($\overline{4-6}$) with numerous axile anatropous ovules with two integuments. *Fruit* a capsule. *Seed* with small embryo in copious endosperm.

Tribe 1. *Asareae*. Herbs with reniform leaves. *Flowers* actinomorphic.

Only genus (perianth campanulate): ☉ **Asarum**.

Tribe 2. *Aristolochieae*. *Flowers* zygomorphic.

Only British genus (perianth tubular, dilated at base): * **Aristolochia**.

Order 8. **Polygonales**. *Leaves* alternate, usually with sheathing bases. *Flowers* haplochlamydeous to heterochlamydeous, actinomorphic. *Ovary* superior, 1-locular, with one basal orthotropous ovule with two integuments.

Fam. POLYGONACEAE. Herbs, rarely shrubs or trees. *Leaves* usually entire with scarious or fleshy amplexicaul, stipular sheath. *Flowers* small in compound inflorescences, cyclic or partially spiral, usually hermaphrodite. *Tepals* 3-6 free. *Stamens* 5-9. $G(3)$ or (2) with as many styles. *Fruit* a nut. *Seed* with copious mealy endosperm. *Embryo* straight or curved.

Subfamily 1. *Rumicoideae*. *Flowers* cyclic. *Endosperm* not fissured.

Tribe *Rumiceae*. *Leaves* with conspicuous sheath.

Stamens in one whorl. *Fruit* not winged. **Rumex**.

Stamens in two whorls. *Fruit* winged.

Rheum (including **Oxyria**).

Subfamily 2. *Polygononoideae*. *Flowers* mostly spiral. *Endosperm* not fissured.

Tribe *Polygoneae*. Herbs, more rarely undershrubs. Only British genus (tepals free, all erect in fruit):

Polygonum (including **Fagopyrum*).

c. (Consisting of the one order **Centrospermae**.) Corolla usually haplochlamydeous, sepaloid or petaloid. Heterochlamydy, however, is not uncommon. (For *d*, Heterochlamydeae, see page 45, for *b* page 37.)

Order 9. **Centrospermae**. Mostly herbs, often with peculiar secondary thickening. *Leaves* often entire, usually exstipulate. *Stipules* scarious if present. *Flowers* homo- or heterochlamydeous. *Stamens* as many as and opposite to perianth segments or $\infty-1$. *Carpels* $\infty-1$, usually united. *Ovary* mostly superior and 1-loc with 1- ∞ campylotropous ovules. *Placentation* basal or free central. Two integuments present. *Embryo* curved. *Seeds* often reniform and with granulate testa. *Perisperm* often present.

Suborder 1. CHENOPODINEAE. *Perianth* homochlamydeous and sepaloid. *Tepals* not more than 5 with *stamens* opposite them. *Ovule* solitary.

Fam. 1. CHENOPODIACEAE. Herbs, rarely shrubs, often with 'bladder-hairs' ('mealiness'). *Leaves* alternate, exstipulate, fleshy, entire or irregularly lobed, sometimes much reduced. *Flowers* small, homochlamydeous. *Tepals* 5, 3, or 1, valvate. *Stamens* as many or fewer opposite the tepals, bent inwards in bud. *Carpels* (2), rarely (3-5). *Ovary* 1-loc with one basal ovule. *Fruit* a nutlet or pyxidium enclosed in persistent perianth.

A. *Cyclolobeae*. *Embryo* annular or horse-shoe shaped.

Tribe 1. *Beteae*. *Flowers* hermaphrodite. *Stamens* 1-5, united below. *Ovary* subinferior. *Stigma* short, broad, papillose within. *Fruit* opening with lid.

Only British genus: **Beta.**

Tribe 2. *Chenopodieae*. *Flowers* mostly hermaphrodite. *Stigma* linear. *Fruit* indehiscent.

Only British genus: **Chenopodium.**

Tribe 3. *Atripliceae*. *Flowers* usually diclinous. ♂ with *perianth* and without *bracteoles*. ♀ without *perianth* and with persistent *bracteoles*.

Only British genus: **Atriplex.**

Tribe 4. *Salicornieae*. *Flowers* hermaphrodite embedded in the leaves.

Only British genus: **Salicornia.**

B. *Spirolobeae*. *Embryo* spirally twisted.

Tribe 5. *Suaedae*. *Leaves* succulent, glabrous. *Bracteoles* small. *Style-arms* papillose all round.

Only British genus: **Suaeda.**

Tribe 6. *Salsoleae*. *Leaves* often with filiform hairs. *Bracteoles* as large or larger than tepals. *Style-arms* papillose within.

Only British genus: **Salsola.**

*Fam. 2. AMARANTACEAE. Very near *Chenopodiaceae* but *perianth* not herbaceous, but scarious and coloured; the *tepals* often end in awns.

Trop. equiv. of Chenopods.

***Amarantus.**

Suborder 2. *PHYTOLACCINEAE. *Flowers* haplo- or heterochlamydeous with a tendency to spiral arrangement of parts. *Stamens* sometimes numerous. *Carpels* sometimes only slightly united.

Fam. 3. *AIZOACEAE. Usually herbs, often with fleshy leaves. *Stipules* absent or scarious. *Flowers* usually haplochlamydeous. *Perianth segments* free or united, sepaloid. *Stamens* 5 or more, the outer ones sometimes represented by petaloid staminodes. *Carpels* (2- ∞). *Ovary* 2- ∞ -loc, rarely 1-loc. *Placentation* basal or free central (parietal in *Mesembryanthemum*).

Tribe *Mesembryanthemeae*. *Ovary* inferior. *Staminodes* present or not.

Staminodes very numerous, petaloid.

**Mesembryanthemum*. (See

Suborder 3. PORTULACINEAE. *Flowers* heterochlamydeous. *Sepals*† 2. *Petals* 4-5.

Fam. 4. PORTULACACEAE. Herbs with entire fleshy leaves and scarious stipules. *Flowers* hermaphrodite, heterochlamydeous. *Sepals* 2. *Petals* 4-5, free or connate below, deciduous. *Stamens* 5- ∞ . *Carpels* (3-5) with 2- ∞ ovules on a basal placenta. *Fruit* capsular.

A. *Stamens* 3-5. *Ovary* superior.

Petals free, *stamens* 5.

**Claytonia*.

Petals connate, *stamens* 3.

Montia.

B. *Stamens* numerous. *Ovary* subinferior.

**Portulaca*.

† These structures may also be regarded as bracteoles. In the N. Am. genus *Lewisia* they are 4-8 in number, subulate, and placed on the scape below the insertion of the perianth.

Suborder 4. CARYOPHYLLINEAE. *Flowers* heterochlamydeous with equal number of sepals and petals, cyclic throughout, sometimes apopetalous.

Fam. 5. CARYOPHYLLACEAE. Herbs, rarely shrubs, with opposite, rarely spiral, entire leaves. *Stipules* 0 or scarious. *Flowers* mostly in dichasia, cyclic, usually heterochlamydeous, 5 rarely 4-merous. *Stamens* usually in two whorls. *Carpels* (5-2) with 1- ∞ ovules on a free central or basal placenta. *Fruit* usually capsular.

Subfamily 1. *Alsinoideae*. *Sepals* free. *Stamens* often perigynous. *Styles* free or united.

a. Petals usually conspicuous. Capsule opening with teeth.

Tribe 1. *Alsineae*. *Stipules* 0. *Styles* free.

Cymes lax. Petals bifid. Capsules globose. **Stellaria**.

Cymes usually lax. Petals bifid. Capsules cylindric.

Cerastium (including **Moenchia**).

Cymes corymbose. Petals jagged. Capsules cylindric.

Holosteum.

Petals entire. Styles 3-4 (carpels fewer than sepals).

Arenaria.

Petals entire or 0. Styles 4-5 (carpels isomerous).

Sagina.

Tribe 2. *Sperguleae*. *Stipules* scarious. *Styles* free.

Petals white. Carpels and styles 5. **Spergula**.

Petals white or pink. Carpels and styles 3.

Spergularia.

Tribe 3. *Polycarpeae*. *Stipules* scarious. *Styles* united below.

Only British genus : **Polycarpon**.

b. Petals inconspicuous or absent. Fruit indehiscent.

Tribe 4. *Paronychieae*. *Stipules* scarious. *Petals* minute, bristle-like. *Ovules* 1-4.

A. Sepals obtuse, flat, green.

Leaves linear, opposite and spiral. Stigmas 3.

Corrigiola.

Leaves ovate-oblong, opposite. Stigmas 2.

Herniaria.

B. Sepals awned, laterally compressed, white.

Illecebrum.

Tribe 5. *Scleranthedeae*. *Stipules* 0. *Petals* usually 0. *Ovules* 1-2.

Only British genus (leaves opposite, connate at base):

Scleranthus.

Subfamily 2. *Silenoideae*. *Sepals* united. *Petals* and *stamens* hypogynous. *Gynophore* often present. *Styles* free.

Tribe 6. *Lychnideae*. *Calyx* with commissural ribs.

A. Fruit a capsule.

Carpels isomerous, alternating with sepals. Petals without appendages. Claw 2-winged. **Agrostemma.**

Carpels oligomerous, or if isomerous opposite sepals. Capsule with septa. Teeth as many as styles.

Viscaria.

As *Viscaria*, but teeth twice as many as styles.

Silene.

Capsule with no septa. Teeth as many as styles.

Lychnis.

As *Lychnis*, but teeth twice as many as styles.

Melandryum.

B. Fruit a berry.

***Cucubalus.**

Tribe 7. *Diantheae*. *Calyx* with no commissural ribs.

Bracts immediately beneath capsule. Petals without appendages. Embryo straight.

Dianthus (including *Tunica*†).

Bracts not immediately beneath capsule. Petals with appendages. Embryo curved. **Saponaria*.

d. **Heterochlamydeae** (Moss). **Orders with predominant heterochlamydy.** (For *c* see page 40.)

a. *Apocarpy* and *Hypogyny* predominant. (For β see page 48.)

Order 10. **Ranales**. *Flowers* spiral, spirocyclic, or cyclic, haplo- to heterochlamydeous, hypogynous to epigynous, actinomorphic or zygomorphic. *Stamens* usually ∞ . *Carpels* $\infty-1$, usually free.

Suborder 1. NYMPHAEINEAE. *Flowers* predominantly spiral. *Ovules* usually scattered over the inner surface of the carpels but sometimes solitary at the apex (*Ceratophyllaceae*).

Fam. 1. NYMPHAEACEAE. Aquatic herbs. *Leaves* floating, often peltate. *Flowers* solitary spiral, spirocyclic or cyclic, homo- or heterochlamydeous, hermaphrodite, actinomorphic. *Flower-axis* convex, or hollow and united to gynaecium. P $6-\infty$, A $6-\infty$, G $3-\infty$ free or united, each with $1-\infty$ ovules on their inner surface. *Ovules* with two integuments. *Seeds* often with aril. *Cotyledons* thick.

Subfamily *Nymphaeoidae*. *Flowers* spirocyclic. *Carpels* ∞ , with ∞ ovules completely covering the septa. *Endosperm* and *perisperm* present.

† An ill-defined genus differing from *Dianthus* in having smaller flowers and broad, membranous sepaline commissures.

Tribe 1. *Nuphareae*. K 6-12, C ∞ . Fruit a berry.
Flowers globose, yellow. **Nuphar.**

Tribe 2. *Tetrasepaleae*. K 4, C ∞ .
Flowers expanded, white, blue, or red. **Nymphaea.**

Fam. 2. CERATOPHYLLACEAE. Submerged aquatics with whorls of transparent, forked leaves, which later become cartilaginous. *Flowers* solitary in the leaf axils, haplochlamydeous, diclinous, monoecious. ♂ *perianth* about 12. *Stamens* 12-16. ♀ *perianth* about 9-10. *Carpel* 1 with one apical, orthotropous ovule with one integument. *Style* long, subulate.

Only genus : **Ceratophyllum.**

Suborder 2. RANUNCULINEAE. *Flowers* spiral to cyclic. *Ovules* on the ventral suture.

Fam. 3. RANUNCULACEAE. Mostly acrid herbs. *Leaves* often divided. *Flowers* rarely completely cyclic, haplo- or heterochlamydeous, in first case with petaloid perianth between which and androecium honey-leaves are often present, hermaphrodite, actinomorphic or zygomorphic. *Stamens* usually numerous, free. *Carpels* ∞ -1 usually free with ∞ -1 ovules. *Fruit* of one-seeded achenes or many-seeded follicles, or a berry (*Actaea*). *Endosperm* copious.

A. *Ovules* on both sides of ventral suture, rarely solitary. *Fruit* a follicle with ∞ -1 seeds.

*Tribe 1. *Paeoniaeae*. *Flowers* usually solitary, without honey-leaves. *Carpels* fleshy. *Outer integument* longer than inner.

Only genus : ***Paeonia.**

Tribe 2. *Helleboreae*. Flowers 1 or several. *Carpel wall* rarely fleshy. *Outer integument* not longer than inner.

I. Honey-leaves without spur. Sometimes absent.

A. Leaves undivided or merely lobed. Honey-leaves absent. **Caltha.**

B. Leaves deeply lobed or compound. Honey-leaves present.

† Leaves palmately lobed or compound.

α. Honey-leaves flat with naked honey-pits.

Trollius.

β. Honey-leaves tubular, at least at base.

Leaves palmately divided, sepals membranous, deciduous. ***Eranthis.**

Leaves pedate. Sepals herbaceous, persistent.

Helleborus.

†† Leaves doubly ternate or pinnate, fruit a berry.

Actaea.

II. Honey-leaves with spur.

1. Flowers actinomorphic with five honey-leaves.

Aquilegia.

2. Flowers zygomorphic with two honey-leaves.

Honey-leaves sessile, dorsal sepal spurred.

***Delphinium.**

Honey-leaves stalked, dorsal sepal hooded.

☉ **Aconitum.**

B. *Ovule* solitary at base of ventral suture. *Achene* one-seeded.

Tribe 3. *Anemoneae* (only tribe).

Shrubs. Leaves opposite. Sepals valvate. **Clematis.**

Herbs. Leaves spiral. Sepals 4-5, inconspicuous. Honey-leaves 0. **Thalictrum.**

Herbs. Leaves spiral. Scape with involucre of 3 leaves. Sepals 4-20. Honey-leaves 0. **Anemone.**

Herbs. Leaves spiral. Sepals 5-8. Honey-leaves 5-16, conspicuous. ***Adonis.**

Herbs. Leaves spiral. Sepals 5, spurred. Honey-leaves small, tubular. Flower-axis ultimately elongate.

Myosurus.

Herbs. Leaves spiral. Sepals 3-5, not spurred. Honey-leaves 5, flat, petaloid. **Ranunculus.**

Fam. 4. BERBERIDACEAE. Herbs, shrubs, or trees. *Leaves* simple or compound. *Flowers* solitary or racemose, cyclic, homo- or heterochlamydeous, hermaphrodite 3-2-merous, actinomorphic. *Perianth* with 2-4 whorls, two whorls of honey-leaves often present. *Stamens* in two whorls, anthers opening by lids or valves. *Carpel* 1, rarely several, with $\infty-1$ ovules on ventral suture or basal. Two integuments. *Fruit* a berry. *Seed* with endosperm.

Subfamily *Berberidoideae*. *Leaves* pinnate, or terminal leaflet alone present. *Honey-leaves* present.

Tribe 1. *Berberideae*. *Inflorescence* on lateral short shoots.

Only British genus (flowers 3-merous, fruit a berry):
Berberis.

*Tribe 2. *Epimediaceae*. *Inflorescence* terminal.

Only British genus (flowers 2-merous. Petals with discolorous appendages, fruit dehiscent):

***Epimedium.**

β . *Syncarpy* and *Hypogyny* dominant. (For α see page 45, for γ see page 54.)

Order II. **Rhoeadales.** Usually herbs. *Flowers* often racemose, cyclic (except androecium in some cases), usually heterochlamydeous, hypogynous, actinomorphic or zygomorphic. *Carpels* ($\infty-2$). *Ovules* with two integuments.

Suborder I. RHOEADINEAE. *Flowers* usually heterochlamydeous. *Sepals* usually 2.

Fam. I. PAPAVERACEAE. Herbs, often with white or coloured latex. *Flowers* hermaphrodite, usually nodding in bud. *Sepals* 2. *Petals* usually 4. *Stamens* ∞ , or if only 4-2, then branched from the base. *Carpels* (2-16) with parietal *placentae* and ∞ *ovules*, or with one basal ovule. *Fruit* usually capsular. *Embryo* minute. *Endosperm* oily.

Subfamily I. *Papaveroideae.* *Petals* without spur. *Stamens* ∞ . *Carpels* 2- ∞ .

Tribe I. *Chelidoniace.* *Latex* yellowish or reddish. *Style* ending in two undivided branches which alternate with the *placentae*. *Stigmatic surface* within and on the edges of these branches.

Only British genus:

☉ **Chelidonium.**

Tribe 2. *Papavereae.* *Latex* yellow or white. *Stigmas* lying above the *placentae*.

a. Fruit long, dehiscent to the base.

Fruit with septum.

Glaucium.

Fruit with no septum. Flowers violet. ***Roemeria.**

b. Fruit elongate, ovoid or globose, dehiscent above.

Stigmas 4-5, style distinct.

Meconopsis.

Stigma sessile, on a disc.

Papaver.

c.

4

Subfamily 2. *Fumarioideae*. *Flowers* mostly transversely zygomorphic. One of two *outer petals* gibbous or spurred. *Stamens* opposite them, each stamen divided from the base into three.

Ovules numerous, fruit a 2-valved capsule.

Corydalis.

Ovules 2, fruit 1-seeded, indehiscent.

Fumaria.

Suborder 2. CAPPARIDINEAE. *Flowers* hetero-chlamydeous. *Sepals* 4 or more.

Fam. 2. CRUCIFERAE. Annual or perennial herbs usually with unicellular simple or branched hairs. *Leaves* spiral, exstipulate. *Flowers* in bractless racemes, hermaphrodite, actinomorphic. K 2 + 2. C 4. A 2 (short) + 4 (long). G (2). *Placentation* parietal, but *ovary* is 2-loc on account of growth of septum (*replum*) from placentae. *Stigmas* 2 on short style above placentae. *Fruit* mostly capsular. *Endosperm* absent.

Position of cotyledons and radicle as follows:

o || notorhizal (radicle incumbent).

o = pleurorhizal (radicle accumbent).

o > > orthoplocous (cotyledons conduplicate).

The following arrangement of the British Cruciferae is according to de Candolle and is adapted from Hooker's *Students' Flora*, ed. 3. See also *Syllabus*, ed. 7, p. 198, where another arrangement is given.

A. Siliquosae. *Pods* usually much longer than broad, dehiscent throughout their whole length, not compressed at right angles to septum.

Tribe 1. *Arabideae* (Siliquosae pleurorhizae). *Seeds* 1-seriate (2-seriate in *Arabis* and *Nasturtium*). *Coty-*

ledons flat. *Radicle* accumbent 0 =. (*Flowers* white, yellow or lilac.)

α. Stigmatic lobes erect, or decurrent to the style.

Matthiola.

β. Stigma small, simple, terminal.

Hairs forked. Lateral sepals pouched. Pods compressed or 4-angled, valves 1-nerved. ***Cheiranthus.**

Hairs simple or 0. Pods terete, valves turgid, not elastic. Seeds minute, 2-seriate. **Nasturtium.**

Pods 4-angled, valves keeled. Seeds oblong. Flowers yellow. **Barbarea.**

Pods flat, valves 1-nerved, not elastic. Flowers mostly white. **Arabis.**

Rhizome not fleshy. Pods flat, valves elastic. Funicle filiform. **Cardamine.**

Rhizome fleshy. Pods flat, valves elastic. Funicle dilated. **Dentaria.**

Tribe 2. *Sisymbrieae* (Siliquosae notorhizae). *Seeds* usually 1-seriate. *Cotyledons* flat. *Radicle* incumbent 0 ||. (*Flowers* as above.)

Glabrous, or hairs simple and spreading. Flowers yellow. Sepals equal. Stigma obtuse. **Sisymbrium.**

Hairs appressed, branched. Flowers yellow. Stigma obtuse. **Erysimum.**

Hairs spreading. Flowers white or lilac, fragrant. Lateral sepals gibbous. Stigmas decurrent on style.

***Hesperis.**

Tribe 3. *Brassiceae* (Siliquosae orthoplocae). *Seeds* 1-2-seriate. *Cotyledons* longitudinally folded 0 > >. *Radicle* incumbent. (*Flowers* yellow.)

Pods terete or angled. Seeds 1-seriate, globose.

Brassica.

Pods compressed. Seeds 2-seriate, compressed.

Diplotaxis.

B. *Siliculosae*. *Pods* short, dehiscent through their whole length.

a. *Latisepatae*. Pods compressed parallel to the replum, which is hence as broad as the pod's greatest diameter.

Tribe 4. *Alysseae* (*Latisepatae pleurorhizae*). *Seeds* 2-seriate. *Radicle* accumbent o =.

Pods oblong, compressed, many-seeded.

Draba (including **Erophila**).

Pods circular, 2-8-seeded.

***Alyssum.**

Pods globose, many-seeded.

Cochlearia.

Tribe 5. *Camelineae* (*Latisepatae notorhizae*). *Seeds* 2-seriate. *Radicle* incumbent o ||.

Tall herbs, flowers small, cauline leaves auricled.

***Camelina.**

Tribe 6. *Subularieae* (*Latisepatae diploclorobae*). As *Camelineae*, but *cotyledons* curved with lengthened bases turned up in direction of radicle.

Scapigerous water-herb with subulate leaves.

Subularia.

b. *Angustiseptae*. *Pod* much compressed at right angles to replum, which is hence narrow.

Tribe 7. *Thlaspeae* (*Angustiseptae pleurorhizae*). *Cotyledons* straight, *radicle* accumbent o =. (*Flowers* white.)

Pods obovate, notched. Petals equal. Filaments without scales.

Thlaspi.

Pods orbicular. Petals white or lilac, very unequal. Filaments without scales.

Iberis.

Pods oblong. Petals white, unequal. Filaments with scales. **Teesdalia.**

Pods oblong. Petals equal. Filaments without scales. **Hutchinsia.**

Tribe 8. *Lepidieae*. *Cotyledons* straight, incurved or longitudinally folded. *Radicle* incumbent o ||. (*Flowers* white.)

Pods many-seeded. **Capsella.**

Pods 2-4-seeded. **Lepidium.**

Tribe 9. *Brachycarpeae* (*Angustiseptae* *diplocolobae*). *Cotyledons* induplicate. *Radicle* incumbent.

Fruit indehiscent, 2-seeded. **Coronopus.**

C. *Nucamentaceae*. *Pods* indehiscent, 1-celled, 1-seeded.

Tribe 10. *Isatideae* (*Nucamentaceae* *notorhizae*). *Radicle* incumbent o ||.

Only British genus: **Isatis.**

D. *Lomentaceae*. *Fruit* with transverse dissepiments, separating into 1-seeded joints.

Tribe 11. *Cakilinae* (*Lomentaceae* *pleurorhizae* o =). *Radicle* accumbent. *Cotyledons* flat.

Pods compressed, of two dissimilar joints.

Only genus: **Cakile.**

Tribe 12. *Raphaneae* (*Lomentaceae* *orthoplocae* o >>). *Radicle* incumbent. *Cotyledons* longitudinally folded.

Pods of a lower slender seedless and an upper globose 1-seeded joint. **Crambe.**

Pods elongate, of 2-several similar joints. **Raphanus.**

Suborder 3. RESEDINEAE. *Flowers* spiro-cyclic, heterochlamydeous.

Fam. 3. RESEDACEAE. Herbs with spiral leaves. *Flowers* in racemes, spiro-cyclic, hermaphrodite, zygomorphic. *Flower-axis* passing into a short gynophore which expands below the stamens into an eccentric or semilunar disc. K 4-8. C 0-8. A 3-10. G (2-6). *Ovary* 1-locular, open above. *Ovules* usually many on parietal placentae. *Fruit* usually a coriaceous capsule open at the top. No endosperm. *Embryo* curved.

Only British genus :

Reseda.

Order 12. **Sarraceniales.** Herbs with spiral leaves adapted to insect catching. *Flowers* spiro-cyclic to cyclic, homo- or heterochlamydeous, hypogynous, actinomorphic. *Carpels* (3-5) with parietal or axile placentation and ∞ ovules. *Seeds* minute with endosperm.

Fam. DROSERACEAE. Herbs, mostly without main root. *Leaves* stipulate, spiral, often rosulate; with digestive glands and irritable hairs or tentacles. *Flowers* cyclic, heterochlamydeous, hermaphrodite, actinomorphic. K 5-4. C 5-4. A 5-4- ∞ . G (5-3). *Styles* 5-3, rarely 1. *Ovary* 1-locular with ∞ -3 parietal or basal ovules. *Capsule* mostly 1-celled, loculicidal. *Seeds* ∞ -3 with endosperm. *Embryo* minute.

Only British genus :

Drosera.

γ. Apocarpy and Hypogyny still occur but Perigyny becomes more frequent. By sinking of the Gynaecium into the hollow flower-axis Syncarpy and Epigynous insertion of the perianth and stamens also takes place. (For β see page 48, for δ see page 61.)

Order 13. **Rosales.** *Flowers* cyclic, rarely spiro-cyclic (*Rosaceae*—*Rosoideae*), usually heterochlamydeous and actinomorphic. *Carpels* often free. *Placentae* closely allied to *Ranunculales*.

sometimes thickened, usually with ∞ ovules.—Most of the families of this order are very difficult to demarcate.

Suborder I. SAXIFRAGINEAE. *Carpels* as many as petals or fewer. *Endosperm* mostly copious (scanty in *Crassulaceae*).

Fam. I. CRASSULACEAE. Mostly succulent herbs. *Leaves* exstipulate. *Flowers* cyclic, heterochlamydeous, 3–30-merous, haplo- or obdiplostemonous, mostly hermaphrodite and actinomorphic. *Petals* free or united. *Carpels* usually isomerous, free or slightly coherent, with glandular scales at base. *Ovules* mostly 2-seriate on the ventral suture. *Fruit* of follicles. *Seeds* small oblong with scanty endosperm.

A. Obdiplostemonous.

* Petals free.

Leaves scattered or 2–3 in a whorl. Flowers 4–5-merous. Sedum.

Leaves rosulate. Flowers 6–20-merous.

*Sempervivum.

** Corolla tubular, 5-lobed.

Cotyledon.

B. Haplostemonous.

Leaves opposite. Flowers 3–5-merous.

Crassula (including Tillaea).

Fam. 2. SAXIFRAGACEAE. Mostly herbs. *Leaves* usually spiral. *Stipules* absent or present as outgrowths from the leaf-sheath. *Flowers* cyclic, usually heterochlamydeous and 5-merous (but *Carpels* mostly oligomerous), hermaphrodite, actinomorphic. *Flower-axis* convex, flat, or concave, in the latter case usually united by its whole length to the carpels. *Stamens* usually

obdiplostemonous, but sometimes haplostemonous, more rarely ∞ . *Carpels* seldom free and as many as petals, mostly fewer, often two. *Styles* mostly free. *Ovary* 2-1-loc with swollen placentae bearing ∞ ovules in several rows. *Seeds* minute with copious endosperm and small embryo.

Subfamily 1. *Saxifragoideae*. Herbs with spiral leaves. *Flowers* mostly 5-merous. *Carpels* 2, rarely 3-4, free or united below. *Ovules* with two integuments.

Tribe 1. *Saxifrageae*. *Carpels* rarely free, but *styles* always free.

Flowers 5-merous with corolla. Placentation axile.

Saxifraga.

Flowers 4-merous with no corolla. Placentation parietal.

Chrysosplenium.

Tribe 2. *Parnassieae*. *Carpels* (3-4) forming syncarpous gynaecium. *Style* short or absent.

Parnassia.

Subfamily 2. *Ribesoideae*. Shrubs with simple exstipulate leaves. *Flowers* in racemes. *Stamens* 5. *Ovary* inferior, 1-loc, with parietal placentae. *Fruit* a berry.

Only genus:

Ribes.

Suborder 2. ROSINEAE. *Carpels* ∞ -1. *Ovules* with two integuments. *Endosperm* scanty or absent.

Fam. 3. PLATANACEAE. Trees with bark separating in large scales. *Leaves* spiral, 3-5-lobed, with large connate *stipules*. *Flowers* in globose heads, diclinous, cyclic, heterochlamydeous, actinomorphic, 3-8-merous, typically isomerous with four alternating whorls but

disturbances of this arrangement by abortion occur. *Stamens* with short filaments and clavate anthers with peltate prolongation of connective. *Carpels* free with 1-2 almost orthotropous ovules with two integuments. *Fruit* a caryopsis. *Endosperm* scanty.

**Platanus*.

Fam. 4. ROSACEAE. Herbs, shrubs or trees. *Leaves* spiral. *Stipules* sometimes adnate to petiole, rarely absent. *Flowers* cyclic, usually heterochlamydeous, actinomorphic, rarely zygomorphic. *Flower-axis* flat, dish- or cup-shaped: sometimes convex in the middle. *Sepals*, *petals* and *stamens* on the edge of the axis, perigynous or epigynous. *Petals* often orbicular and concave. *Stamens* mostly 2-4 times as many as sepals, or ∞ , rarely 1-5, bent inwards in bud. *Carpels* as many as sepals or 2-3 times as many or ∞ , rarely 1-4, free or united with the inner wall of the flower-axis, 1-locular with usually two anatropous ovules. *Styles* apical or on ventral side of carpels. *Fruit* follicular or indehiscent, or drupaceous or a false fruit by union of carpels with flower-axis. *Endosperm* scanty or absent. *Cotyledons* mostly fleshy, plano-convex.

Subfamily **Spiraeoideae*. *Filaments* narrowed upwards from broad base. *Carpels* 12-1 whorled, rarely sunk into flower-axis or on gynophore. *Ovules* ∞ -2. *Fruit* usually of follicles.

*Tribe *Spiraeae*. Shrubs rarely herbs. *Seeds* not winged.

**Spiraea*†.

† The plants often known in Britain as *S. Ulmaria* and *S. Filipendula* belong more properly to the genus *Ulmaria*, subfamily *Rosoideae*, tribe *Ulmarieae*.

Subfamily *Pomoideae*. *Stipules* distinct. *Carpels* 5-2, united with the inner wall of the hollow flower-axis, usually syncarpous (i.e. united to each other). *False fruit* formed of carpels + axis + lower part of calyx.

A. Carpels free on their ventral sides (i.e. the sides to which the styles are attached).

Fruit a drupe with 3-5 partially exserted stones.

Cotoneaster.

B. Carpels united.

Inner wall of carpel cartilaginous when ripe. **Pirus.**

Inner wall of ripe carpel hard, fruit hence a drupe.

Leaves lobed. Flowers in corymbose cymes.

Crataegus.

Leaves undivided. Flowers solitary.

***Mespilus.**

Subfamily *Rosoideae*. *Carpels* either ∞ , on conical gynophore, rarely few not enclosed in axis, or 1- ∞ enclosed in persistent flower-axis: each carpel with 1-2 ovules. *Fruit* always indehiscent.

Tribe I. *Potentilleae*. *Flower-axis* flat. *Stamens* ∞ , narrowed upwards from broad base. *Carpels* mostly ∞ , on convex gynophore.

No epicalyx. Carpels each with two ovules. Fruit drupaceous.

Rubus.

Epicalyx present. Carpels with one ovule. Fruit dry, indehiscent. Style not persistent. Achenes on large fleshy receptacle.

Fragaria.

As *Fragaria* but achenes on small dry receptacle.

Potentilla.

Epicalyx. Fruit indehiscent with persistent style.

Leaves simple. Scape 1-flowered. Petals 8-9.

Dryas.

Leaves pinnate. Flowers several. Petals 5. **Geum**.

Tribe 2. *Ulmarieae*. *Flower-axis* flat or slightly concave. *Filaments* narrow at base, soon falling.

Only genus: **Ulmaria**.

Tribe 3. *Sanguisorbeae*. *Flower-axis* pitcher-shaped, enclosing and usually hardening around two or more achenes.

Epicalyx of bristles. Petals present. Styles terminal. **Agrimonia**.

Epicalyx of four leaves. No petals. Styles basal or ventral. **Alchemilla**.

Epicalyx and petals both absent. **Poterium**.

Tribe 4. *Roseae*. *Flower-axis* tubular, enclosing ∞ carpels, becoming soft when ripe.

Only genus: **Rosa**.

Subfamily *Prunoideae*. *Carpel* 1, rarely more. *Style* terminal. *Ovules* 2, pendulous. *Fruit* drupaceous.

Only British genus: **Prunus**.

Fam. 5. LEGUMINOSAE. Herbs, shrubs or trees. *Leaves* of spiral stipulate, often compound. *Flowers* usually in racemes, cyclic, heterochlamydeous, 5-merous, hypogynous, usually hermaphrodite and zygomorphic. *Stamens* usually 10, 9 or all often united. *Carpel* mostly solitary with usually ∞ ovules on the ventral suture which is directed backwards. *Style* terminal. *Fruit* a pod or lomentum. *Endosperm* scanty or absent.

Subfamily *Papilionatae*. *Roots* in symbiosis with *Bacillus radiculicola*. *Flowers* zygomorphic. *Corolla* with descending imbrication in bud.

Tribe 1. *Genisteae*. Shrubs or herbs with simple or palmate leaves. *Leaflets* entire. *Calyx* 2-lipped. All ten *stamens* usually united. *Pod* 2-valved.

Herbs. Leaves palmate. *Calyx* deeply 2-lipped. *Pod* coriaceous with spongy septa. ***Lupinus**.

Shrubs. Leaves 1-foliolate. *Calyx* with two short deeply toothed lips. **Genista**.

Shrubs. Mature leaves spinescent or scaly. *Calyx* deeply 2-lipped, coloured. **Ulex**.

Shrubs. Leaves 1-3-foliolate or absent. *Calyx* with two short minutely toothed lips. **Cytisus**.

Tribe 2. *Trifolieae*. Herbs. *Leaves* usually pinnately 3-foliolate. *Veins of leaflets* ending in teeth. *Upper stamen* usually free.

A. All ten *stamens* united. Keel beaked. **Ononis**.

B. Upper *stamen* separate. Keel obtuse.

a. Corolla deciduous, claws of petals free.

I. Flowers in heads, umbels, or short racemes.

Pod straight or curved, mostly linear, often beaked.

Trigonella.

Pod usually spiral, sometimes broadly oval and bent.

Medicago.

II. Flowers in long racemes. *Pods* obovoid.

Melilotus.

b. Corolla usually persistent, claws of all or of four lower petals united to the staminal tube. **Trifolium**.

Tribe 3. *Loteae*. Herbs. *Leaves* pinnate with entire leaflets. *Upper stamen* separate or united. *Pod* 2-valved without longitudinal septum.

Pod enclosed by the calyx.

Anthyllis.

Pod long, exserted.

Lotus.

Tribe 4. *Astragaleae*. Herbs. *Leaves* pinnate: *leaflets* entire. *Upper stamen* usually free. *Pod* 2-valved with longitudinal septum derived from suture.

Keel obtuse. Pod-septum from dorsal suture.

Astragalus.

Keel beaked or tip incurved. Pod-septum from ventral suture.

Oxytropis.

Tribe 5. *Hedysareae*. *Upper stamen* usually separate. *Fruit* indehiscent, either a lomentum of several 1-seeded joints or consisting of only one such joint.

Keel obliquely truncate. Fruit 1-2-seeded, not jointed.

Onobrychis.

Keel beaked. Fruit jointed, curved, laterally compressed.

Hippocrepis.

Keel obtuse. Fruit terete, many-jointed.

Ornithopus.

Keel beaked. Fruit terete or 4-angled, jointed.

***Coronilla.**

Tribe 6. *Vicieae*. Mostly herbs. *Leaves* pinnate with no terminal leaflet but instead a tendril or point. *Cotyledons* thick.

Style filiform, hairy below or all round.

Vicia.

Style flat, hairy above.

Lathyrus.

8. *The flowers have predominantly 5 or 4 whorls. Apocarpy and Isomery still occur, but Syncarpy and Oligomery of the Gynaecium preponderate. Pleiometry of Gynaecium rare. (For γ see page 54.)*

Order 14. **Geraniales**. Herbs (the British species): *Flowers* cyclic, heterochlamydeous, usually 5-merous.

Handwritten: *cp. Sapindales*

Carpels (5-2) often separating from each other when ripe. *Ovules* usually 2-1, more rarely ∞ , anatropous, pendulous, with ventral raphe and micropyle directed upwards, or, when more than one ovule is present, single ones sometimes occur with dorsal raphe and micropyle directed downwards.

Suborder I. GERANIINEAE. *Flowers* heterochlamydeous, mostly actinomorphic. Usually obdiplostemonous (i.e. *stamens* twice as many as petals, and *carpels* when isomerous opposite petals). Haplostemony more rarely, or individual stamens may abort in zygomorphic flowers. *Anthers* opening by longitudinal slits. *Carpels* isomerous or oligomerous. *Ovules* with two integuments.

Fam. I. GERANIACEAE. Herbs with lobed or divided, often stipulate *leaves*. *Flowers* 5-merous, actinomorphic. No real disc present. *Petals* imbricate or convolute in bud. *Stamens* 10, sometimes only five fertile. *Carpels* usually 5, each with 1-2 ovules. *Capsule* splitting into five beaked mericarps. *Seeds* with endosperm.

Tribe I. *Geranieae*. *Mericarps* separating by means of elastic awns.

Fertile stamens 10. Styles glabrous within, twisting in flat spiral after dehiscence. **Geranium†.**

Fertile stamens 5. Styles silky within, twisting in helicoid spiral after dehiscence. **Erodium.**

† The so-called *Geraniums* of cultivation belong to the South African genus *Pelargonium*, which differs from the other *Geranieae* in having a sepaline spur which is adnate to the pedicel and best seen in section. The flowers are slightly zygomorphic. The fruit resembles that of *Erodium*.

Fam. 2. OXALIDACEAE. Perennial herbs. *Leaves* often ternate, showing sleep-movements. *Flowers* 5-merous. No real *disc*. *Stamens* 10, united below. *Carpels* isomerous. *Fruit* a capsule or berry, loculicidal if dehiscent. *Endosperm* fleshy.

Only British genus:

Oxalis.

Fam. 3. LINACEAE. Herbs with simple, entire *leaves*. *Flowers* 4-5-merous, actinomorphic. *Petals* often convolute in bud. No real *disc*. *Stamens* 5-20, united in a ring below. *Gynaecium* quite syncarpous. *Ovary* often with extra, imperfect dissepiments from dorsal sutures. *Fruit* a capsule or drupe.

Tribe *Eulineae*. *Stamens* in one whorl. *Fruit* a capsule.

Flowers 5-merous. *Sepals* entire.

Linum.

Flowers 4-merous. *Sepals* with 2-3 teeth. **Radiola.**

Suborder 2. POLYGALINEAE. *Flowers* actinomorphic or zygomorphic. *Stamens* in two whorls. *Anthers* opening by pores. *Carpels* (2), median.

Fam. 4. POLYGALACEAE. Herbs or shrubs. *Leaves* usually simple, entire, exstipulate. *Flowers* 5-merous, zygomorphic. *Sepals* 5, of which two are petaloid and wing-like. *Petals* by abortion only 3. *Stamens* 8. *Carpels* with one rarely with 2-4 ovules. *Fruit* a capsule or drupe.

Only British genus:

Polygala.

Suborder 3. TRICOCCEAE. *Flowers* actinomorphic, diclinous, often very reduced. *Carpels* usually (3). *Seed* mostly with caruncle.

Fam. 5. EUPHORBIACEAE. Herbs, shrubs or trees, often with latex. *Leaves* usually spiral, often stipulate. *Inflorescence* mostly compound. *Stamens* as many as

sepals or twice as many or ∞ or 1. *Carpels* usually (3). *Fruit* usually a capsule splitting into three mericarps. *Endosperm* copious.

Subfamily *Crotonoideae*. Each *carpel* with one *ovule*.

Tribe 1. *Acalypheae*. *Latex* absent. *Calyx* valvate. *Flowers* racemose. ♂ usually without corolla.

Only British genus: **Mercurialis.**

Tribe 2. *Euphorbieae*. *Latex* present. *Flowers* in *cyathia*†. ♂ without corolla and usually without calyx, and with only one stamen.

Only British genus: **Euphorbia.**

Suborder 4. CALLITRICHINEAE. Characters of the family. (Systematic position doubtful. Possible affinities with Verbenaceae.)

Fam. 6. CALLITRICHACEAE. Slender glabrous herbs, often submerged. *Leaves* opposite entire, upper often rosulate and floating. *Flowers* monoecious, naked, often with two falcate bracteoles. ♂ of one terminal *stamen*. ♀ of two transverse *carpels*. A longitudinal septum makes ovary 4-loc. *Fruit* of four drupaceous mericarps. *Seeds* with endosperm.

Only genus: **Callitriche.**

Order 15. **Sapindales.** Mostly woody plants. (Impatiens is the only British herbaceous genus.) Characters of Geraniales but ovules in the reverse position, either pendulous with dorsal raphe and micropyle directed upwards, or ascending with ventral raphe and micropyle directed downwards.

is of *carpel*.† A *Cyathium* is a condensed cymose inflorescence resembling a single flower.

pp. *Geraniales*.

Suborder 1. BUXINEAE. *Flowers* haplochlamydeous. *Ovules* with two integuments.

Fam. 1. BUXACEAE. Woody plants. *Leaves* entire, evergreen, exstipulate. *Flowers* diclinous, sometimes with rudiments of the aborted sporophylls, actinomorphic. *Stamens* 4- ∞ . *Carpels* usually (3) each with 2-1 ovules. *Styles* separate. *Fruit* a loculicidal capsule or drupe. *Seeds* with endosperm.

Only British genus (leaves opposite): **Buxus.**

Suborder 2. EMPETRINEAE. *Flowers* heterochlamydeous. Each *carpel* with one ascending ovule with one integument. *Carpels* not separating when ripe.

Fam. 2. EMPETRACEAE. Ericoid shrubs with linear exstipulate leaves deeply furrowed beneath. *Flowers* diclinous with rudiments of the aborted sporophylls, actinomorphic. K, C and A 2-3. G (2-9). *Fruit* a drupe.

Only British genus (leaves spiral, stamens 3):

Empetrum.

Suborder 3. CELASTRINEAE. *Flowers* heterochlamydeous, always actinomorphic, diplostemonous or haplostemonous. *Gynaecium* tending towards oligomery, rarely isomalous.

Fam. 3. AQUIFOLIACEAE. Trees with spiral, mostly evergreen, simple leaves. *Flowers* 4-5-merous, dioecious, actinomorphic. *Disc* absent. *Petals* often connate at base and adnate to the isomalous stamens. *Carpels* (4-6) each with 1-2 pendulous ovules with one integument. *Fruit* a drupe.

Only British genus (petals connate at base): **Ilex.**

Fam. 4. CELASTRACEAE. Trees or shrubs with simple opposite or spiral leaves. *Stipules* absent or

deciduous. *Flowers* small, often green, 4-5-merous, mostly hermaphrodite. *Disc* usually conspicuous. *Petals* imbricate. *Stamens* 4-5, inserted on the disc. *Carpels* (2-5) each with $\infty-1$ ovules. *Fruit* a capsule or berry. *Seeds* often arillate.

Only British genus (leaves opposite, carpels isomerous): **Euonymus.**

*Fam. 5. STAPHYLEACEAE. Shrubs or trees. *Leaves* usually opposite and pinnate, often with stipules and stipels. *Flowers* in panicles, 5-merous. *Stamens* 5, inserted outside the disc. *Carpels* (2-3), free above. *Ovules* ∞ -few. *Fruit* capsular. *Embryo* straight. *Endosperm* copious, fleshy.

Capsule inflated.

***Staphylea.**

Suborder 4. SAPINDINEAE. *Flowers* heterochlamydeous, typically diplostemonous but some stamens and carpels abort, actinomorphic or obliquely zygomorphic. *Ovules* with two integuments.

Fam. 6. ACERACEAE. Trees. *Leaves* opposite, simple, lobed or pinnate, exstipulate. *Flowers* actinomorphic, axis flat or concave. K and C 4-10. A 4-10, usually 8. G (2) each with two almost orthotropous ovules. *Fruit* indehiscent, winged, usually 1-seeded. No *endosperm*.

Only British genus (fruit wings lateral): **Acer.**

*Fam. 7. HIPPOCASTANACEAE. Trees. *Leaves* opposite, palmate, exstipulate. *Flowers* in conspicuous, primarily racemose inflorescences, obliquely zygomorphic. K 5. C 4-5. A 5-8. G (3) each with two ovules. *Capsule* 1-seeded, 2-3-valved. No *endosperm*.

Only British genus (calyx tube long): ***Aesculus.**

Suborder 5. BALSAMININEAE. *Flowers* heterochlamydeous, zygomorphic, haplostemonous, with united anthers.

Fam. 8. BALSAMINACEAE. Herbs or shrubs. *Leaves* opposite or alternate. *Flowers* always hermaphrodite. *Sepals* usually 3 (2 anterior, if present, minute), posterior spurred. *Petals* primitively 5, *anterior* large, external in bud, *lateral* and *posterior* connate on each side so that there are apparently only three petals. *Stamens* 5, filaments short, anthers cohering. *Carpels* (5) each with ∞ ovules. *Capsule* loculicidal, opening elastically, rarely indehiscent. *Seeds* ∞ . *Endosperm* absent.

Only British genus (capsule opening elastically):

Impatiens.

Order 16. **Rhamnales.** *Flowers* cyclic, diplochlamydeous, sometimes apopetalous. *Stamens* in one whorl opposite the petals. *Carpels* (5-2) each with 1-2 ascending ovules, with dorsal, lateral or ventral raphe and two integuments.

Fam. RHAMNACEAE. *Stems* usually woody, rarely herbaceous, often climbing. *Leaves* simple, often 3-5-nerved. *Stipules* small. *Flowers* small, greenish or yellowish, often in axillary cymes, 5-4-merous, perigynous or epigynous. *Calyx lobes* triangular, valvate. *Petals* small, often clawed and hooded and enclosing the stamens, or absent. *Stamens* inserted on calyx tube at edge of disc. *Carpels* (5-2). *Fruit* various, often a drupe. *Seeds* mostly with endosperm.

Tribe *Rhamneae*. *Serial buds* not present (as in *Colletia*). *Ovary* superior or inferior.

Only British genus (disc lining the calyx tube):

Rhamnus.

Order 17. **Malvales.** *Flowers* cyclic (*androecium* not always so), heterochlamydeous, hermaphrodite, rarely

zygomorphic. *Calyx* and *corolla* usually 5-merous, *sepals* mostly valvate in bud. *Stamens* ∞ or in two whorls, the inner of which is branched. *Carpels* (2- ∞) each with 1- ∞ anatropous ovules with two integuments.

Suborder MALVINEAE. *Sepals* mostly valvate: *mucilage* present.

Fam. 1. TILIACEAE. Mostly trees with entire or toothed stipulate *leaves*. *Mucous-tubes* present in pith and bark. *Flowers* usually hermaphrodite. K and C 5-merous. *Calyx* valvate. *Corolla* sometimes absent. *Stamens* usually ∞ inserted on disc free or in 5-10 bundles. *Anthers* with two thecae. *Carpels* (2- ∞) each with 1- ∞ ovules. *Style* 1. *Fruit* 2- ∞ -locular or 1-locular by abortion. *Endosperm* mostly present.

Only British genus (wing-like bract adnate to peduncle): **Tilia.**

Fam. 2. MALVACEAE. Herbs, shrubs or trees with simple or lobed stipulate *leaves*. *Sepals* 5, valvate, often with *epicalyx*. *Petals* 5, adnate to base of staminal column, twisted in bud. *Stamens* usually ∞ , *filaments* forming a tube. *Anthers* with one theca. *Pollen grains* large, spiny. *Carpels* (5- ∞) each with 1- ∞ ovules. *Styles* as many or twice as many as *carpels*. *Fruit* a capsule or schizocarp. *Seeds* often woolly. *Endosperm* scanty or absent.

Tribe *Malveae*. *Style-arms* as many as *carpels*. *Fruit* a schizocarp.

Epicalyx of 3-6 connate segments. Axis longer than fruits. **Lavatera.**

Epicalyx of 6-9 connate segments. Axis shorter than fruits. **Althaea.**

Epicalyx of 3 distinct segments. **Malva.**

ε. *Flowers spirocyclic or of 5-4 whorls. Syncarpy the rule (Apocarpy in the more primitive, exotic families). Tendency for gynaecium to become sunk in flower-axis.* (For δ see page 61, for ζ see page 71.)

Order 18. **Parietales.** *Flowers spirocyclic or cyclic often with indefinite number of stamens and carpels, heterochlamydeous, rarely apopetalous, hypogynous or epigynous. Carpels more or less united often with parietal placentae, which, however, may meet in the middle. Ovules very seldom basal.* Pariet
wall
(Har

The suborders of Parietales may stand in phylogenetic relationship to some of the earlier orders, particularly Ranales and Rhoadales. Suborder Flacourtiineae (see page 71) shows possible affinities with Cucurbitaceae. Yet the Cucurbitaceae possess so many important peculiarities that they cannot have been derived directly from the Flacourtiineae.

Suborder 1. THEINEAE†. *Stamens often ∞. Gynaecium free, on convex or flat flower-axis. Placentation often axile. Endosperm containing oil and proteid grains.*

Fam. 1. GUTTIFERAE. *Trees or shrubs, more rarely herbs (most British species are herbaceous). Leaves opposite, entire, usually exstipulate, often with pellucid glands. K and C often 5, sepals imbricate. Stamens often ∞ and united in bundles. Carpels (3-5) with ∞-1 ovules with two integuments. Placentation parietal or axile. Seeds without endosperm.*

Subfamily *Hypericoideae.* *Stamens ∞ in 2-5 bundles. Styles usually free. Fruit a capsule, berry or drupe.*

Tribe *Hypericeae.* *Herbs and shrubs. Ovary often 3-5-loc. Capsule septicidal.*

Only British genus (petals 5, unequal-sided) :

Hypericum.

† Cf. *Guttiferales* of Bentham and Hooker.

Suborder 2. TAMARICINEAE. *Stamens* in whorls or if ∞ in bundles. *Gynaecium* free on flat flower-axis. *Endosperm* starchy or absent.

Fam. 2. ELATINACEAE. Usually small plants growing in wet places and rooting at the nodes. *Leaves* opposite or whorled, stipulate. *Flowers* small, cyclic, 2-5-merous. *Sepals* and *petals* each 2-5, imbricate. *Stamens* in one or two whorls. *Ovary* 2-5-loc with ∞ axile ovules with two integuments. *Styles* as many as loculi, free. *Capsule* septicidal.

Only British genus (leaves spathulate): **Elatine.**

Fam. 3. FRANKENIACEAE. Usually small shrubs. *Stems* jointed at the nodes. *Leaves* opposite, exstipulate. *Flowers* 4-6-merous. *Calyx* gamosepalous. *Petals* free, with ligular appendages. *Stamens* 4- ∞ . *Carpels* 4-2 with ∞ ascending ovules on parietal placentae. *Style* slender. *Stigma* 2-5-lobed. *Capsule* dehiscing between placentae. *Seeds* with endosperm.

Only British genus (petals 5, usually with appendages): **Frankenia.**

*Fam. 4. TAMARICACEAE. Herbs or shrubs with minute alternate, entire leaves. *Flowers* mostly 4-5-merous. *Stamens* as many as petals or twice as many or ∞ in groups. *Carpels* (5-2). *Ovary* usually 1-loc with ∞ basal or parietal ovules. *Fruit* a capsule. *Seeds* hairy.

Tribe *Tamariceae*. *Flowers* racemose. *Seeds* with apical tuft of hairs.

Only British genus (stamens free): ***Tamarix.**

Suborder 3. CISTINEAE. *Stamens* ∞ , not in bundles. *Gynaecium* free on flat or convex axis. *Endosperm* with starch.

Fam. 5. CISTACEAE. Herbs and shrubs. *Leaves* mostly opposite. Stellate and glandular hairs, the latter with essential oils, often present. $K\ 5-3$. $C\ 5-3$ or 0 . $A\ \infty$. $G\ (5-10)$. *Ovary* usually 1-celled with parietal placentae with ∞ or 2 more or less orthotropous ovules. *Capsule* dehiscing between placentae. *Seeds* with endosperm.

Only British genus (ovules ∞ , capsule 3-valved):

Helianthemum.

Suborder 4. FLACOURTIINEAE. *Stamens* often 5. *Gynaecium* free on convex or tubular axis, rarely adnate to axis. *Endosperm* copious containing oil and proteid bodies.

Fam. 6. VIOLACEAE. Herbs or shrubs with alternate stipulate leaves. *Flowers* 5-merous (except gynaecium), actinomorphic or zygomorphic. *Petals* sometimes united. *Stamens* 5. *Carpels* (3) each with $1-\infty$ anatropous ovules with two integuments on parietal placentae. *Fruit* a loculicidal capsule or berry.

Tribe *Violeae*. *Corolla* zygomorphic, *two anterior stamens* spurred.

Only British genus (flowers solitary in the leaf axils):

Viola.

§. *The flowers are cyclic and the sinking of the gynaecium into the hollow flower-axis is general: connation of gynaecium and flower-axis predominates.* (For ϵ see page 69.)

Order 19. **Myrtiflorae.** Herbs, shrubs, or trees, bundles often bicollateral. *Flowers* cyclic, heterochlamydeous, rarely apopetalous, haplo- or diplostemonous, usually actinomorphic. *Flower-axis* concave. *Gynaecium* syncarpous and mostly united to axis.

Suborder 1. THYMELAEINEAE. Usually shrubs with entire leaves. *Flower-axis* more or less tubular

(except in ♂ Elaeagnaceae). *Gynaecium* of 2-4 carpels, free from the flower-axis.

Fam. 1. THYMELAEACEAE. Shrubs or trees with acrid juice and tenacious, reticulate bast. *Leaves* entire, alternate or opposite, exstipulate. *Flowers* hermaphrodite, heterochlamydeous or apopetalous, diplo- or haplostemonous, 5-4-merous. *G* usually 1 with one pendulous ovule. *Style* 1, terminal or lateral. *Fruit* often a drupe.

Subfamily *Thymelaeoideae*. *Flowers* diplostemonous or haplostemonous. *Petals* scale-like or absent. *Carpel* 1 with one ovule.

Only British genus (filaments and style very short):

Daphne.

Fam. 2. ELAEAGNACEAE. Shrubs or trees with silvery or brown scales. *Flower-axis* in ♂ flowers flat, in ♀ and ♂ flowers tubular. *Flowers* mostly 4-merous, homochlamydeous. *Androecium* diplostemonous in ♂ flowers; in ♀ haplostemonous and with stamens opposite perianth lobes. *Carpel* 1 with one basal ovule. *Fruit* a nut enclosed in the fleshy axis.

Only British genus (sepals 2, stamens 4): **Hippophaë.**

Suborder 2. MYRTINEAE. Herbs or shrubs. *Leaves* more often opposite. *Flowers* with tubular axis and 2-∞ carpels forming syncarpous gynaecium which is usually united to axis. *Ovules* with one integument.

Fam. 3. LYTHRACEAE. Herbs or shrubs. *Leaves* opposite or whorled, entire. *Flowers* heterochlamydeous or apopetalous, usually 4-6-merous, hermaphrodite, actinomorphic or zygomorphic. *Flower-axis* cup-shaped or tubular. *Calyx* valvate with intersepalar stipules. *Petals* on edge of flower-axis, sometimes absent. *Stamens*

twice as many as sepals or $1-\infty$, inserted deeper than the petals. *Carpels* (2-6) each with $\infty-2$ ovules, free from the *flower-axis*. *Ovary* 2-6-loc, *ovules* ∞ , axile. *Style* 1 with capitate stigma. *Fruit* a capsule. *Seeds* without endosperm.

Tribe *Lythraeae*. *Septa of ovary* imperfect upwards.

Flower-axis tubular. Corolla large. Stamens 12.
Fruit dehiscent. **Lythrum.**

Flower-axis campanulate. Corolla small or absent.
Stamens 6. Fruit indehiscent. **Peplis.**

Fam. 4. EPILOBIACEAE (Onagraceae). Usually herbs. *Leaves* alternate or opposite, exstipulate. *Flowers* heterochlamydeous, usually hermaphrodite and actinomorphic. *Flower-axis* tubular. K 2-4, P 2-4, rarely absent. A 4-8, sometimes partially staminodal. G usually (4), united to axis, each with $\infty-1$ ovules. *Style* 1. *Stigma* entire or 4-lobed. *Fruit* various, usually with many seeds. *Endosperm* scanty or absent.

Tribe *Oenotheraeae*. *Ovary* quite inferior.

Calyx persistent. Petals short or absent. Stamens 4.
Capsule short. **Ludwigia.**

Calyx deciduous. Petals 4, pink or purple. Stamens 8. Capsule long. Seeds with hairs. **Epilobium.**

Calyx deciduous. Petals 4 (yellow in Brit. sp.).
Capsule long. Seeds with membranous margin.

***Oenothera.**

Calyx deciduous. Petals 2, white. Stamens 2. Fruit dry, indehiscent. **Circaea.**

Shrubs. Flowers pendulous. Calyx petaloid. Fruit a berry. ***Fuchsia.**

Fam. 5. HALORRHAGACEAE. Herbs of very diverse habit. *Flowers* small, heterochlamydeous, or often apopetalous. *Stamens* twice as many as *sepals* or fewer. *Anthers* long, 4-angled. *Carpels* usually (4) united to the tubular axis, each with one pendulous ovule. *Styles* separate. *Fruit* a nut or drupe. *Seeds* with endosperm.

Subfamily 1. *Halorrhagoideae*. *Petals* 2-4 or 0. *Carpels* (2-4) each with one ovule.

Only British genus (leaves whorled, deeply pinatifid. *Petals* 2-4): **Myriophyllum**.

Suborder 3. HIPPURIDINEAE. *Flowers* epigynous. *Stamen* 1. *One carpel* with one ovule without integument.

Fam. 6. HIPPURIDACEAE. Aquatic herbs. *Leaves* linear, in whorls. *Flowers* minute, axillary, naked. *Carpel* 1 with one undivided style, stigmatic throughout its length.

Only genus (stems simple, erect): **Hippuris**.

Order 20. **Umbellales (Umbelliflorae)**. *Flowers* mostly in umbels, cyclic, heterochlamydeous, usually haplostemonous, epigynous, 4-5-merous. *Carpels* (5-1), each carpel with one (rarely two) pendulous anatropous ovules with one integument. *Seeds* with copious endosperm.

Fam. 1. ARALIACEAE. Mostly trees and shrubs, often twining or climbing by aid of roots. *Leaves* alternate, simple or compound, often stipulate. *Flowers* mostly in umbels or heads, 5-merous. *Calyx* often with obscure limb. *Stamens* usually 5. *Carpels* (1-∞) each

usually with one ovule. *Micropyle* facing outwards. *Fruit* a berry or drupe.

Tribe *Schefflereae*. *Petals* valvate. *Endosperm* sometimes lobulate.

Only British genus (leaves simple, endosperm lobulate): **Hedera.**

Fam. 2. UMBELLIFERAE. Herbs with hollow internodes. *Leaves* alternate with conspicuous *sheath*, and usually compound *lamina*. *Flowers* in simple or compound umbels, 5-merous, haplostemonous. *Calyx* usually inconspicuous. *Petals* epigynous, tips often inflexed. *Stamens* at base of stylopodium. *Anthers* versatile. G (2), median. *Styles* 2 arising from epigynous disc (*stylopodium*). *Fruit* separating into two *mericarps* pendulous on the *carpophore*, each *mericarp* with five *primary ridges*, i.e. two lateral next to the commissure and three dorsal. Four *secondary ridges* are sometimes present between the primary. Oil canals (*vittae*) occur in the grooves between the *primary ridges*: two or more are present on the commissural face of the *mericarp*. *Seed* adherent to pericarp. *Endosperm* copious. *Embryo* minute.

Subfamily 1. *Hydrocotyloideae*. *Leaves* often simple (peltate in the only British sp.). *Flowers* in heads or simple umbels. *Fruit* with woody fibrous *endocarp*. *Carpophore* undivided. *Vittae* absent or in the primary ridges: never in the furrows.

Tribe 1. *Hydrocotyleae*. *Fruit* laterally compressed. (*Commissure* narrow.)

Only British genus:

Hydrocotyle.

Subfamily 2. *Saniculoideae*. Leaves various. Inflorescence as above. Endocarp soft. Styles long surrounded at base by annular stylopodium. Stigmas capitate. Vittae various.

Tribe 2. *Saniculeae*. Commissure broad.

Leaves palmately divided. Umbels irregularly compound. Fruit globose, with hooked bristles. Ridges obscure. **Sanicula.**

Leaves palmate. Umbels simple. Bracts large, coloured. Fruit ovoid. Ridges wrinkled or toothed. ***Astrantia.**

Leaves with spinous teeth. Flowers capitate. Fruit rough, ridges absent. **Eryngium.**

Subfamily 3. *Apioideae*. Leaves various. Flowers usually in compound umbels. Endocarp soft (there may be layer of *stereom* beneath *epidermis*). Style arising from apex of stylopodium. Vittae at first in the furrows, later variously arranged.

A. *Haplozygieae*. Primary ridges more conspicuous than the secondary. Vittae usually obvious in the furrows.

α. Seed furrowed ventrally by the raphe.

Tribe 3. *Scandicineae*. Crystal-glands present around carpophore.

Subtribe 1. *Scandicinae*. Fruit long-cylindrical and beaked, smooth or with short spines.

i. Vittae several in each furrow. **Conopodium.**

ii. Vittae 1 in each furrow, or absent.

Fruit smooth, over 2 cm. long, with prominent obtuse ridges and long beak. **Scandix.**

Fruit slightly rough, over 2 cm. long. Ridges very acute. Carpophore split to middle. **Myrrhis.**

Fruit rather rough, 5 mm. long. Ridges vanishing upwards. Carpophore undivided or shortly bifid.

Chaerophyllum.

Fruit smooth, under 1 cm. long. Beak short, ribbed.

Anthriscus.

Subtribe 2. *Caucalinae*. Fruit ovoid, secondary ridges spinous.

Only British genus: **Caucalis** (including **Torilis**).

*Tribe 4. *Coriandreae*. Crystal-glands absent around carpophore. Fruit with woody layers beneath the epidermis. *Mericarps* coherent.

***Coriandrum.**

Tribe 5. *Smyrnieae*. No crystal-glands. Fruit ovoid. *Commissure* narrow. *Mericarps* separating.

One vitta in each furrow; ridges entire, slender.

Physospermum.

Several vittae in each furrow. Stylopodium flattened. Ridges crenate.

Conium.

Several vittae in each furrow. Stylopodium conical. Ridges obscure.

Smyrnium.

β . Seed flat ventrally. *Raphe* often projecting towards carpophore.

Tribe 6. *Ammineae*. Dorsal and commissural primary ridges all alike. Seed semicircular in section.

Subtribe 1. *Carinae*. Ridges not prominent. *Commissure* narrow.

1. Petals entire, with acute or inflexed tips. Vittae 1-3 in each furrow.

Leaves simple, entire. Flowers yellow. **Bupleurum.**

Leaves compound. Flowers white, dioecious. ♂
flowers with narrower petals. **Trinia.**

Leaves compound. Flowers white, hermaphrodite.
Apium.

2. Petals 2-lobed. Tip long, inflexed. One vitta
in each furrow.

Calyx teeth minute or absent. Vittae as long as
fruit. **Carum.**

Calyx teeth minute or absent. Vittae only in upper
half of fruit. **Sison.**

Calyx teeth leafy, ovate, acute. Vittae long. **Cicuta.**

3. Petals as 1, but vittae usually several in each
furrow.

Leaves pinnate. Calyx teeth acute. Ridges obtuse.
Sium.

Leaves twice 3-nate. Calyx teeth absent. Ridges
slender. Vittae 0. **Aegopodium.**

Calyx teeth absent. Vittae many. **Pimpinella.**

Subtribe 2. *Seselinae*. Ridges prominent, sometimes
winged, lateral ones forming continuation of commissural
face of fruit.

1. Fruit almost cylindrical. Ridges not thickened
or corky.

Calyx teeth minute. Petals white, notched. **Seseli.**

Calyx teeth 0. Petals yellow, entire. **Foeniculum.**

2. Fruit as above. Primary ridges acute; outer coat
of pericarp loose, corky. **Crithmum.**

3. Fruit as above. Primary ridges thick, lateral
ridges forming a corky rim round the carpel.

Bracteoles short, whorled.

Oenanthe.

Bracteoles long, unilateral.

Aethusa.

4. Fruit as above, lateral ridges thickened or winged.

Silaus.

5. Fruit dorsally compressed. Primary ridges broad, thick.

Seed grooved ventrally, vittae several.

Meum.

Seed flat or slightly concave ventrally, vittae many or obscure.

Ligusticum.

Seed biconvex: one vitta in each of the dorsal furrows.

Selinum.

Tribe 7. *Peucedaneae*. *Lateral ridges* winged, much broader than the three dorsal ridges which are often only feebly developed. Seed narrow in section.

Subtribe 1. *Angelicinae*. *Lateral ridges* of opposite carpels not appressed but gaping.

Only British genus:

Angelica (including **Archangelica*).

Subtribe 2. *Ferulinae*. *Lateral ridges* appressed, forming a wing round the fruit.

Only British genus:

Peucedanum.

Subtribe 3. *Tordyliinae*. As *Ferulinae* but wing hardened.

Vittae clavate. Margins of wing thin.

Heracleum.

Vittae slender. Margins of wing thick.

Tordylium.

B. *Diplozygieae*. *Secondary ridges* as large as or larger than the primary.

Tribe 8. *Dauceae*. *Secondary ridges* spinous.

Daucus.

Fam. 3. CORNACEAE. Trees or shrubs. *Leaves* opposite or spiral, usually entire, exstipulate. *Flowers* small, in cymes, umbels, or heads, 4-5-merous, mostly haplostemonous. *Carpels* (4-1) with an epigynous disc, each carpel with one ovule. *Microphyll* facing outwards or inwards. *Style* 1. *Fruit* a drupe or berry.

Subfamily *Cornoideae*. *Ovary* inferior. *Raphe* dorsal.

Only British genus (leaves opposite. Petals 4, valvate): **Cornus**.

Subclass 2. **METACHLAMYDEAE** (Symptetales). *Perianth* in advanced stage of development, always originally double, and the inner whorl gamopetalous. Polypetalous forms occur but they are closely related to gamopetalous forms.

Stamens 2. **Polypetaly**, as well as **sympetaly**, occurs. **Two whorls or one whorl of stamens**. **Hypogyny predominates**, but **epigyny** also occurs. (For B see page 85.)

Order 1. **Ericales**. Shrubs and herbs, rarely trees, with simple *leaves*. *Flowers* 4-5-merous, obdiplostemonous, or the antipetalous whorl of *stamens* not developed. *Petals* free or united. *Stamens* hypogynous or epigynous, more rarely united at base with the *petals*. *Carpels* (2- ∞), when isomerous opposite petals. *Ovary* superior or inferior. *Ovules* with one *integument*.

Fam. 1. PIROLACEAE. Evergreen or colourless perennial herbs with spiral *leaves*. *Flowers* solitary or racemose, 5-4-merous, obdiplostemonous. *Petals* free or united. *Stamens* hypogynous. *Carpels* (5-4) each with ∞ minute *ovules* on fleshy *placentae*. *Capsule*

loculicidal. *Seeds* with loose *testa* and fleshy *endosperm*. *Embryo* of few cells without cotyledons.

Subfamily 1. *Pirolloideae*. *Anthers* recurved before flowering, then erect, opening by two *pores*. *Pollen* in tetrads.

Only British genus (herbs with broad evergreen leaves): *Pirola* (including *Moneses*).

Subfamily 2. *Monotropoideae*. *Anthers* always erect at the apex of the filament, opening by slits. *Pollen* simple.

Tribe *Monotropeae*. *Ovary* 4-5-loc below, 1-loc above.

Only British genus (colourless root-parasites):

***Monotropa*.**

Fam. 2. ERICACEAE. Usually low shrubs. *Leaves* mostly evergreen. *Flowers* solitary or racemose, 5-4-merous, obdiplostemonous. *Corolla* usually sympetalous. *Stamens* on the edge of a hypogynous or epigynous disc. *Anther-thecae* free, tubular, often spreading above and with basal or dorsal *appendages*. *Pollen* in tetrads. *Gynaecium* syncarpous. *Ovules* 1-∞ in each carpel on axile *placentae*. *Style* 1, *stigma* capitate. *Fruit* various. *Seeds* with loose *testa* and copious *endosperm*. *Embryo* often very short.

A. *Septicidal capsule*. *Petals* free or united. *Anthers* without appendages. *Testa* loose, often winged.

Subfamily 1. *Rhododendroideae*. As above. *Buds* scaly.

*Tribe 1. *Ledaeae*. *Flowers* actinomorphic. *Petals* free. *Seeds* with long wings.

(Petals 5, spreading.)

**Ledum*.

*Tribe 2. *Rhododendreae*. *Corolla* sympetalous, often weakly zygomorphic. *Seeds* surrounded by broad wing. (Stamens exserted.)

**Rhododendron* (including **Azalea*).

Tribe 3. *Phyllodoceae*. *Petals* usually united. *Seeds* globose or 3-angled, not winged. *Embryo* cylindrical.

Corolla campanulate. *Stamens* 5. *Anthers* opening by slits. *Capsule* 2-3-valved. *Loiseluria*.

Corolla urceolate. *Stamens* 10. *Anthers* opening by pores. *Capsule* 5-valved. *Phyllodoce*.

Corolla urceolate. *Stamens* 8. *Anthers* opening by pores. *Capsule* 4-valved. *Dabeocia*.

B. *Berry*, *drupe* or *loculicidal capsule*. *Petals* united, deciduous. *Anthers* appendaged or prolonged into tubes.

Subfamily 2. *Arbutoideae*. *Ovary* superior.

Tribe 4. *Andromedaeae*. *Buds* scaly. *Capsule* loculicidal.

Only British genus (*Corolla* pitcher-shaped):

Andromeda.

Tribe 5. *Arbuteae*. *Buds* naked. *Berry* or *drupe*. *Anthers* with two long reflexed appendages.

Berry many-seeded with firm endocarp. *Arbutus*.

Drupe with several 1-seeded stones. *Arctostaphylos*.

Subfamily 3. *Vaccinioideae*. *Ovary* inferior.

Tribe 6. *Vaccinieae*. *Petals* united. *Stamens* epigynous. *Ovary* sharply demarcated from peduncle.

Only British genus (*Filaments* straight, *Fruit* a berry):

Vaccinium (including *Oxycoccus*).

C. *Capsule*. *Petals* united, persistent, scarious. *Anther-thecae* opening by pores above. *Appendage* usually on back of connective.

Subfamily 4. *Ericoideae*.

Tribe 7. *Ericaceae*. *Buds* naked. *Leaves* small. *Carpels* with many ovules. *Capsule* many-seeded.

Calyx small, green. Corolla with four lobes. Capsule loculicidal. *Erica*.

Calyx sepaloid, exceeding corolla. Corolla deeply divided. Capsule septicidal. *Calluna*.

Order 2. **Primulales**. *Flowers* 5 (rarely 4-8)-merous, usually actinomorphic. *Corolla* nearly always gamopetalous. *Stamens* in one whorl, inserted on corolla opposite the lobes (*antipetalous*). *Ovary* 1-loc with $\infty-1$ ovules which have two *integuments* on free central or basal *placenta*. *Style* . 1

Fam. PRIMULACEAE. Mostly perennial herbs. *Leaves* usually spiral, exstipulate. *Stamens* opposite corolla lobes. *Ovary* usually superior with ∞ whorled or spirally arranged ovules on a free central *placenta*. (*bulb*) *Style* 1. *Capsule* with usually many-angled and faceted *seeds* sunk in cavities on the *placenta*.

Tribe 1. *Androsaceae*. *Corolla* imbricate or quin-cuncial in bud, *lobes* eventually erect or spreading, never reflexed (in British genera).

Subtribe 1. *Primulinae*. Land plants. *Corolla* lobes entire or bifid. *Capsule* with valves.

Only British genus (*Leaves* broad. *Flowers* conspicuous. *Corolla-tube* long): *Primula*.

Subtribe 2. *Hottoniinae*. Water plants. Corolla lobes entire. Capsule with valves.

Only genus (Leaves all submerged, pinnately divided):
Hottonia.

*Tribe 2. *Cyclamineae*. Rootstock often tuberous. Corolla lobes reflexed.

Only British genus (Scapes 1-flowered): ***Cyclamen**.

Tribe 3. *Lysimachieae*. Corolla tube short. Lobes twisted in bud, never reflexed.

Subtribe 1. *Lysimachiinae*. Capsule with valves.

Leaves subrosulate towards top of scape. Corolla white, lobes 5-9. **Trientalis**.

Leaves spiral, opposite or whorled. Corolla yellow, lobes 5. **Lysimachia**.

Subtribe 2. *Anagallidinae*. Capsule opening by lid. Corolla absent. Calyx campanulate, coloured. **Glaux**. Corolla shorter than calyx. Filaments glabrous. **Centunculus**.

Corolla longer than calyx. Filaments villous.

Anagallis.

Tribe 4. *Samoleae*. Corolla quincuncial in bud. Ovary half inferior.

Only genus (Bracts displaced on to the pedicels):

Samolus.

Order 3. **Plumbaginales**. Corolla polypetalous or gamopetalous. Stamens in one whorl opposite the petals. Gynaecium (5). Styles 5 or style 5-fid. Ovary 1-loc with one basal anatropous ovule with long funicle and two integuments. (Perhaps connected with *Centrospermae*.) as *Primulales* may be, too).

Fam. PLUMBAGINACEAE. Often maritime herbs, rarely shrubs. *Leaves* entire, spiral, often radical. *Inflorescence* mostly compound. *Flowers* with bracts and bracteoles. *Calyx* often scarious and coloured. *Petals* free or united. *Fruit* enclosed in calyx.

Tribe *Staticeae*. *Inflorescence* compounded of cincinni. *Stamens* united to the *corolla*. *Styles* united only at base.

Cymes very dense. Styles hairy.	Armeria.
Cymes lax. Styles glabrous.	Statice.

B. **Sympetaly dominant.** **Stamens** always in one whorl. **Union of Carpels** sometimes imperfect. **Hypogyny** usual. (For A see p. 80, for C see p. 87.)

Order 4. **Contortae.** Shrubs and herbs, rarely trees. *Leaves* usually opposite, entire and exstipulate. *Flowers* mostly 5-merous, more rarely 2-6-merous, mostly sympetalous. *Corolla* usually twisted in bud. *Stamens* usually isomerous (two in *Oleaceae*) and inserted on the *corolla*, rarely hypogynous. *Carpels* 2, sometimes free below.

Suborder I. OLEINEAE. *Stamens* 2. *Ovules* with one integument.

Fam. I. OLEACEAE. Woody plants, sometimes climbing, rarely herbs. *Leaves* opposite or whorled, simple or pinnate. *Flowers* racemose or cymose, 2-6-merous, mostly sympetalous, more rarely with free petals or apopetalous, hermaphrodite or diclinous. *Corolla* 4, 5, or 6, free or united. *Stamens* 2, *filaments* short. *Carpels* (2), each *carpel* with usually two anatropous ovules. *Fruit* a capsule, berry or drupe.

Subfamily *Oleoideae*. *Ovules* pendulous from apex of loculi.

Tribe 1. *Fraxineae*. *Petals* free or united only at base, sometimes absent. *Fruit* winged.

Only British genus (Fruit-wing distally produced):

Fraxinus.


*Tribe 2. *Syringae*. *Corolla* sympetalous. *Fruit* a loculicidal capsule with winged *seeds*.

***Syringa.**

Tribe 3. *Oleeae*. *Petals* 4, free or united. *Fruit* a berry or drupe, mostly 1-seeded.

Only British genus:

Ligustrum.

Suborder 2. GENTIANINEAE. *Stamens* as many as petals. *Ovary* superior, 1-2-loc with ∞ ovules which have one integument on parietal or axile *placentae*. 

Fam. 2. GENTIANACEAE. Annual or perennial, usually glabrous, bitter herbs, rarely shrubs. *Leaves* opposite, entire, exstipulate. *Flowers* cymose, usually 4-5-merous, sympetalous, hermaphrodite, actinomorphic. *Sepals* free or united. *Corolla* mostly twisted in bud. *Stamens* as many as petals, inserted on the corolla. *Carpels* (2) with usually ∞ ovules. *Ovary* mostly 1-loc with parietal *placentae*. *Fruit* a 2-valved capsule. *No* (

Subfamily 1. *Gentianoideae*. *Leaves* opposite. *Corolla* convolute or imbricate in bud. *Pollen grains* not compressed.

Tribe 1. *Chironieae*. *Style* slender. *Placentae* projecting.

Leaves connate. *Corolla* large, rotate. *Stamens* 6-8.

Blackstonia (= Chlora auct.).

Corolla minute, funnel-shaped. Stamens 4. Stigma peltate. **Microcala.**

Corolla salver-shaped. Stamens 4. Stigmas with two lamellae. **Cicendia.**

Corolla funnel-shaped, persistent. Stamens 5. Anthers twisted. **Centaureium.**

Tribe 2. *Swertiaeae*. Style short. Placentae not projecting.

Only British genus (Corolla subclavate): **Gentiana.**

Subfamily 2. *Menyanthoideae*. Leaves alternate. Corolla induplicate valvate. Pollen grains compressed.

Leaves 3-foliate. Corolla funnel-shaped.

Menyanthes.

Leaves orbicular, floating. Corolla rotate. **Villarsia.**

Fam. 3. APOCYNACEAE. Shrubs or trees, rarely entirely herbaceous, often climbing. Latex present. *Leaves* opposite, quite entire. *Corolla* usually twisted in bud. *Stamens* 4-5, *anthers* basifixed. *Carpels* usually 2, free below. *Styles* united, ending in discoid or globose head which bears the stigmas. *Fruit* various.

Subfamily *Plumerioideae*. Stamens quite free or loosely attached to stigmatiferous head. *Thecae* mostly without *appendages*. *Seeds* usually without tuft of hairs.

Tribe *Plumerieae*. *Carpels* 2, free below the style.

Only British genus (disc of 2 scales, filaments clavate):

Vinca.

C. **Sympetaly constant.** Only one whorl of stamens present. *Carpels* usually 2, always perfectly united. **Zygomorphy frequent.** (For B see page 85.)

a. *Perianth hypogynous.* (For b see page 98.)

Order 5. **Tubiflorae**. Mostly herbs. *Flowers* typically with four isomeric whorls, but the *gynaecium* usually shows reduction. When zygomorphic the *androecium* too shows reduction. *Stamens* epipetalous. *Ovules* with one integument.

Suborder 1. CONVOLVULINEAE. *Leaves* mostly spiral. *Flowers* usually actinomorphic. *Carpels* with few, often only two, ovules, *micropile* facing downwards. *Fruit* seldom separating into *nutlets* (never in British genera).

Fam. 1. CONVOLVULACEAE. Often left-handed twiners. *Flowers* often conspicuous, 5-4-merous. *Corolla* mostly induplicate valvate in bud. *Stamens* inserted at base of corolla-tube. *Carpels* (2), each with two basal erect ovules. *Styles* separate or united. *Fruit* usually capsular.

Subfamily 1. *Convolvuloideae*. Autotrophic, with green *foliage leaves*. *Corolla* without scales.

Tribe *Convolvuleae*. *Style* 1. *Capsule* dehiscent, 4-seeded.

Bracteoles small. Stigma slender. **Convolvulus**.
Bracteoles large, enclosing calyx. Stigma broad.

Calystegia.

Subfamily 2. *Cuscutoidae*. Leafless parasites. *Corolla* usually with ring of *scales* beneath the *stamens*.

Only genus : **Cuscuta**.

Fam. 2. POLEMONIACEAE. Herbs, rarely shrubs. *Leaves* spiral, exstipulate, sometimes pinnate. *Flowers* 5-merous, usually actinomorphic. *Corolla* usually twisted to right in bud. *Carpels* (3), each with usually ∞ *ovules*. *Style* 1, trifid. *Capsule* mostly loculicidal.

Subfamily *Polemonioideae*. Herbs. *Embryo* green, with ovate or linear cotyledons.

Tribe *Polemonieae*. *Flowers* actinomorphic.

Only British genus (Leaves pinnate. Corolla almost rotate): **Polemonium.**

Suborder 2. BORRAGININEAE. As *Convolvulineae*, but *micropile* facing upwards. *Fruit* often separating into *nutlets*.

Fam. 3. BORRAGINACEAE. Mostly hispid herbs. *Leaves* alternate, undivided. *Flowers* in cincinni, 5-merous, mostly zygomorphic. *Corolla* often with *hollow folds* opposite the lobes. *Stamens* with short *filaments* and often subulate *anthers*. *Carpels* (2), each with two anatropous *ovules*. *Ovary* usually 4-loc by false septa, 4-lobed. *Style* 1, arising from between the lobes. *Stigma* simple or bifid. *Fruit* usually dividing into four nutlets.

Subfamily *Borraginoideae*. *Ovary* deeply lobed. *Fruit* of four or fewer 1-seeded nutlets. *Endosperm* absent.

Tribe 1. *Cynoglosseae*. *Nutlets* inserted by broad ventral surfaces on to conical axis, their tips not projecting beyond point of attachment.

Corolla rotate. Nutlets glabrous with inrolled borders. ***Omphalodes.**

Corolla funnel-shaped. Nutlets with hooked bristles. **Cynoglossum.**

*Tribe 2. *Eritrichieae*. Tips of *nutlets* projecting beyond narrow point of insertion. Axis conical.

Calyx with alternating teeth. Nutlets tubercled.

***Asperugo.**

Calyx without alternating teeth. Nutlets with bristly rim.

***Lappula.**

Tribe 3. *Anchuseae*. *Axis* nearly flat. *Nutlets* with concave surface of insertion which is often surrounded by a ring.

a. Corolla with hollow scales.

Corolla tubular, 5-toothed, hollow scales linear, anthers included, without appendages.

Symphytum.

Corolla rotate, hollow scales short. Filaments with dorsal appendages. Anthers exserted, conniving to form a cone.

Borrigo.

Corolla salver-shaped, hollow scales short. Stamens included, with no appendages.

Anchusa (including **Lycopsis**).

b. Corolla without hollow scales. **Pulmonaria.**

Tribe 4. *Lithospermeae*. *Nutlets* erect with small flat surface of insertion.

No calyx-tube. Corolla imbricate in bud. Stamens included. Nutlets stony.

Lithospermum.

Calyx-tube short. Corolla imbricate in bud. Stamens protruding. Nutlets soft.

Mertensia.

Calyx-tube long. Corolla twisted in bud. Nutlets smooth.

Myosotis.

Tribe 5. *Echieae*. *Flowers* zygomorphic. *Corolla* usually without scales.

Only genus (Calyx regular):

Echium.

Suborder 3. VERBENINEAE. *Leaves* mostly opposite or whorled. *Flowers* mostly zygomorphic. Each *carpel*

with two ovules, rarely with only one. *Fruit* often of nutlets.

Fam. 4. VERBENACEAE. Herbs, shrubs or trees. *Leaves* mostly opposite. *Flowers* mostly zygomorphic. *Calyx* tubular. *Corolla* often with curved *tube* and 2-lipped *limb*. *Stamens* mostly 4, didynamous. *Carpels* mostly (2), each with two *ovules*; *micropyle* facing downwards. *Ovary* usually ultimately 4-loc. *Style* terminal, simple. *Fruit* mostly drupaceous (nutlets in British species).

Tribe *Verbeneae*. Mostly herbs. *Flowers* in spikes or racemes. *Loculi* of ovary divided or not. No *endosperm*.

Only British genus :

Verbena.

Fam. 5. LABIATAE. Aromatic herbs and shrubs. *Leaves* and *branches* opposite or whorled. *Flowers* usually zygomorphic, in cymes often forming *false whorls*. K 5. C 5, *limb* often 2-lipped. A 4, didynamous, or two *stamens* and two *staminodes*, or two *stamens* and no *staminodes*, 5th *stamen* rarely represented by *staminode*. G (2), each with two erect anatropous *ovules*. The *carpels* are folded in between the *ovules*. *Fruit* dividing into four *nutlets*, or fewer by abortion.

Subfamily 1. *Ajugoideae*. *Calyx-tube* 10-11-ribbed. *Corolla* 1 or 2-lipped, rarely almost actinomorphic. *Stamens* 4, rarely 2. *Ovary* usually shortly divided, or to one-third its length. *Nutlets* obovoid with dry *pericarp*. *Endosperm* very scanty. *Embryo* straight.

Tribe 1. *Ajugeae*. *Calyx* 10-ribbed. *Corolla* either almost actinomorphic or (as in British genera) with very

short *upper lip* and *lower lip* reflexed. *Stamens* 4 or 2. *Anthers* 2-celled. *Nutlets* warty or reticulate.

Corolla persistent, with ring of hairs within. Upper lip 2-lobed. **Ajuga.**

Corolla deciduous, without ring of hairs. Upper lip absent. **Teucrium.**

Subfamily 2. *Scutellarioideae*. *Calyx* 2-lipped. *Corolla* 2-lipped, with helmet-shaped upper lip. *Stamens* 4. *Ovary* 4-lobed. *Nutlets* more or less globular, with dry pericarp, attached by small basal insertion to gynophore-like torus. *Seed* transverse. *Endosperm* absent. *Embryo* bent.

(Calyx-tube dilated above to form hollow pouch.)

Scutellaria.

Subfamily 3. *Stachydoideae*. *Calyx-tube* with 5–15 ribs. *Corolla* almost actinomorphic or markedly zygomorphic. *Stamens* 4 or 2. *Ovary* 4-lobed. *Nutlets* ovoid, obovoid or tetrahedric, with dry pericarp and small basal attachment. *Seed* straight. *Embryo* straight.

Tribe 2. *Marrubieae*. *Calyx* tubular or campanulate, 5–10-toothed. *Corolla* usually hidden by *calyx*. *Stamens* and *style* included.

Only British genus:

Marrubium.

Tribe 3. *Nepeteae*. *Calyx* 15-ribbed. *Corolla* 2-lipped. *Stamens* 4, posterior (inner) pair longer or alone present.

Only British genus:

Nepeta.

Tribe 4. *Stachyeeae*. *Calyx* 5–10-ribbed. *Upper lip* of *corolla* concave. *Stamens* 4, parallel, ascending under the upper lip, *anterior* (outer) pair longer.

Subtribe 1. *Brunellinae*. *Calyx* 2-lipped. Lips closing together in fruit. *Upper lip of corolla* concave.

Only British genus : **Brunella.**

Subtribe 2. *Melittinae*. *Calyx* broadly campanulate not closing in fruit. *Corolla-tube* dilated throughout.

Only British genus : **Melittis.**

Subtribe 3. *Lamiinae*. *Calyx* tubular not closing in fruit. *Corolla-tube* not dilated below.

A. Anther-thecae transverse during dehiscence. *Corolla* with hollow scales on palate. **Galeopsis.**

B. Dehiscing anther-thecae vertical or oblique.

1. Nutlets sharply 3-angular with obtuse apex.

Leaves serrate. *Calyx* teeth not spinous. Throat of corolla dilated. **Lamium.**

Leaves incised. *Calyx* teeth spinous. Throat of corolla not dilated. ***Leonurus.**

2. Nutlets ovoid with rounded apex.

Calyx funnel-shaped, teeth dilated at base or united to form a ring. **Ballota.**

Calyx tubular or campanulate, teeth 3-angular, not dilated or united. **Stachys.**

Tribe 5. *Salviae*. *Calyx* campanulate or tubular. *Corolla* with helmet-shaped or sickle-shaped upper lip. *Stamens* 2 (the anterior pair). *Connective* linear, bearing one fertile theca.

Only British genus : **Salvia.**

Tribe 6. *Satureieae*. *Calyx* with five equal teeth or 2-lipped. *Corolla* with flat lobes, almost actinomorphic or 2-lipped. *Stamens* 4 or 2, equal or the anterior pair longer.

1. Corolla distinctly 2-lipped. Stamens 4, ascending under upper lip, spreading above.

Corolla-tube recurved. Upper lip concave. **Melissa.**

Corolla-tube almost straight. Upper lip flat.

Satureia (*Calamintha* auct.).

2. Corolla distinctly 2-lipped. Stamens 4, spreading.

Erect. Leaves broad. Calyx tubular. **Origanum.**

Procumbent. Leaves small. Calyx 2-lipped.

Thymus.

3. Corolla almost actinomorphic. Stamens spreading.

Fertile stamens 4.

Mentha.

Two anterior stamens alone fertile.

Lycopus.

Suborder 4. SOLANINEAE. *Flowers* actinomorphic, or more often zygomorphic, typically 5-merous. *Stamens* 5 or 4 or 2. *Carpels* rarely (5), mostly (2), usually with ∞ *ovules*. *Fruit* mostly capsular, never dehiscent quite to the base.

a. Vascular bundles bicollateral. (For β see p. 95.)

Fam. 6. SOLANACEAE. Herbs or shrubs with spiral *leaves*. *Flowers* terminal, single and often supra-axillary, or in cymose, often supra-axillary, *inflorescences*, mostly 5-merous, hermaphrodite, actinomorphic, or rarely zygomorphic. *Corolla-lobes* mostly folded in bud. *Stamens* 5 (in zygomorphic flowers one may be staminodal). *Carpels* (2), oblique. *Ovules* ∞ on the septum. *Style* 1. *Fruit* a berry or capsule. *Seeds* with endosperm.

Tribe 1. *Solaneae*. *Ovary* 2-locular.

Shrubs. Corolla salver-shaped. Stamens attached above middle of corolla-tube. Fruit a berry. ***Lycium.**

Herbs. Corolla campanulate. Stamens attached at base of tube. Fruit a berry. **Atropa.**

Herbs. Corolla funnel-shaped. Capsule opening by lid. **Hyoscyamus.**

Herbs or shrubs. Corolla rotate. Anthers conniving. Fruit a berry. **Solanum.**

*Tribe 2. *Datureae*. Ovary 4-locular by growth of secondary dissepiments.

(Capsule with spines.) ***Datura.**

β . Bundles collateral. (For α see p. 94.)

I. Ovary 2-locular, usually with ∞ axile ovules. (For II see p. 97.)

Fam. 7. SCROPHULARIACEAE. Herbs, shrubs or trees. Leaves spiral or opposite. Inflorescence various, flowers never terminal. Flowers 5-merous, hermaphrodite, more or less zygomorphic. Calyx usually persistent. Stamens rarely 5, mostly 4 or 2. G (2), median (not oblique), each with ∞ -few anatropous ovules on the dissepiment. Style 1. Fruit a capsule or berry.

Subfamily 1. *Pseudosolanoideae*. Leaves mostly spiral. The two posterior (upper) corolla-lobes, or the upper lip, cover the lateral lobes in bud. Five fertile stamens present.

Tribe 1. *Verbasceae*. Corolla-tube short or absent.

Only British genus (Corolla rotate. Stamens 5):

Verbascum.

Subfamily 2. *Antirrhinoideae*. Lower leaves at least opposite. Aestivation as in *Pseudosolanoideae*. Fifth (posterior) stamen staminodal or absent.

Tribe 2. *Antirrhineae*. Flowers axillary or racemose, zygomorphic. Corolla-tube distinct, spurred or saccate at base. Capsule usually opening by pores.

1. Corolla-tube saccate at base. **Antirrhinum.**
2. Corolla-tube spurred at base.
Leaves narrow, pinnately veined. Flowers in terminal racemes. Capsule with valves. **Linaria.**
Leaves narrow, pinnately veined. Flowers solitary. Capsule with pores. **Elatinoides.**
Leaves cordate, digitately nerved and lobed. Capsule with small trifold valves. ***Cymbalaria.**

Tribe 3. *Cheloneae*. Flowers ultimately cymose, zygomorphic. Corolla-tube not spurred or saccate.

Only British genus: **Scrophularia.**

Tribe 4. *Gratioleae*. Inflorescence racemose or flowers axillary. Corolla-tube not spurred or saccate.

Leaves opposite. Flowers large. Stigma 2-lamellate. ***Mimulus.**

Leaves fascicled. Flowers minute. Stigma clavate. **Limosella.**

Subfamily 3. *Rhinanthoideae*. The two posterior (upper) corolla-lobes or the upper lip are covered in bud by the lateral corolla-lobes.

a. Corolla-lobes all nearly flat.

Tribe 5. *Digitaleae*. Not parasitic. Anther-cells united at end.

Corolla-tube short. Stamens 4-8, equal.

Sibthorpia.

Corolla-tube short. Stamens 2, equal. **Veronica.**

Large herbs. Corolla-tube long, dilated. Stamens 4, didynamous. **Digitalis.**

Small herbs. Corolla-tube cylindric. Stamens 4, didynamous. ***Erinus.**

b. The two upper *corolla-lobes* form a helmet-shaped upper lip.

Tribe 6. *Rhinantheae*. Partial and entire parasites.

1. Leaves opposite, narrow. Seeds 1-4, large, not winged. **Melampyrum.**

2. Leaves opposite. Seeds ∞ , minute, not winged.

Upper lip of corolla almost entire. **Bartschia.**

Upper lip of corolla 2-lobed. **Euphrasia.**

3. Leaves opposite. Seeds few, winged.

Rhinanthus.

4. Leaves alternate. Seeds ∞ , minute, not winged.

Pedicularis.

II. *Ovary* 1-locular with ∞ *ovules* on more or less parietal *placentae*. (For I see p. 95, for III see below.)

Fam. 8. OROBANCHACEAE. Annual and perennial, usually brownish, parasitic herbs. *Leaves* scale-like. *Flowers* usually racemose, 5-merous, zygomorphic. *Corolla* distinctly 2-lipped, tube curved. *Stamens* 4, didynamous. *Carpels* (2), median. Each *carpel* with two parietal placentae with ∞ ovules. *Style* 1. *Fruit* a loculicidal capsule.

Subterranean leaves fleshy and with cavities. Flowers racemed. Stamens shortly exserted. **Lathraea.**

Subterranean leaves membranous or o. Flowers sub-spikate. Stamens included. **Orobanche.**

III. *Ovary* usually 1-celled with free central *placenta* bearing ∞ *ovules*.

Fam. 9. LENTIBULARIACEAE. Herbs, mostly aquatic, or growing in wet places. *Roots* often absent. *Leaves*

radical or whorled, undivided or multifid. *Flowers* 5-merous, mostly zygomorphic. *Corolla* usually distinctly 2-lipped. *Stamens* usually 2. *Carpels* (2), median. *Ovules* many on the free central *placenta*. *Fruit* capsular.

Tribe *Utricularieae*. *Corolla* zygomorphic with distinct *spur*.

Land plants. Leaves radical, entire. Corolla throat open. **Pinguicula.**

Aquatic plants. Leaves floating, multifid. Corolla throat closed. **Utricularia.**

Order 6. **Plantaginales.** Mostly herbs. *Leaves* usually spiral. *Flowers* actinomorphic, 4-merous (except *gynaecium*). *Corolla* scarious. One whorl of *stamens*. *Carpels* usually 2.

Fam. PLANTAGINACEAE. Scapigerous herbs. *Leaves* mostly radical, parallel-veined. *Flowers* small, anemophilous, in spikes. K 4, persistent. C 4, scarious, *lobes* spreading. A 4, inserted on corolla-tube (hypogynous in *Litorella*). *Anthers* large, versatile. *Carpels* usually (2). *Ovary* 2-loc or 4-loc by spurious septum. *Ovules* usually several, anatropous, peltate, on the septum. *Fruit* a capsule dehiscing transversely or a nut. *Seeds* with endosperm.

Land plants. Flowers many, spiked, hermaphrodite. Capsule dehiscing transversely. **Plantago.**

Water plant. Flowers few, monoecious. Fruit a nut. **Litorella.**

b. Perianth epigynous. (For *a* see p. 87.)

a. Stamens free. (For β see page 102.)

Order 7. **Rubiales.** Herbs, shrubs and trees. *Leaves* opposite, usually undivided. *Flowers* 5-4-merous, isomerous or *androecium* and *gynaecium* oligomerous, usually actinomorphic. *Ovary* inferior with one or several loculi, each with $\infty-1$ anatropous *ovules*.

A. *Stamens* as many as corolla-lobes. (For B see page 101.)

Fam. 1. RUBIACEAE. Herbs, shrubs or trees, with opposite entire *leaves*. *Stipules* always present, interpetiolar (foliaceous in British species). *Flowers* 5-4-merous, usually actinomorphic. *Sepals* mostly valvate. *Carpels* mostly (2), more rarely (1- ∞), each carpel with 1- ∞ anatropous *ovules*. *Style* 1, with capitate or divided *stigma*. *Fruit* various.

Subfamily *Coffeoideae*. *Carpels* each with one *ovule*.

Tribe *Galieae*. Herbs. *Leaves* and *foliaceous stipules* forming whorls. *Ovules* attached to septum.

A. Sepals distinct, more than half as long as ovary. Corolla funnel-shaped with long tube. **Sherardia.**

B. Sepals minute or absent.

a. Corolla funnel-shaped. **Asperula.**

b. Corolla rotate.

Flowers often 5-merous. Fruit succulent. **Rubia.**

Flowers 4-merous. Fruit dry. **Galium.**

Fam. 2. CAPRIFOLIACEAE. Usually shrubs or small trees. *Leaves* opposite, usually exstipulate. *Stipules* when present (as in *Sambucus*) not interpetiolar. *Flowers* actinomorphic or zygomorphic. *Carpels* (2-5), each with

1- ∞ axile, pendulous *ovules*. *Style* simple or divided. *Fruit* a berry or drupe, rarely a capsule.

Tribe 1. *Sambuceae*. *Leaves* pinnately divided. *Anthers* opening outwards. *Style* very short. *Fruit* a drupe. (Very near *Valerianaceae*.)

Only British genus:

Sambucus.

Tribe 2. *Viburneae*. *Leaves* entire or lobed. *Anthers* opening inwards. *Carpels* (1-5), each with one *ovule*. *Style* usually very short. *Fruit* a berry or drupe.

Only British genus:

Viburnum.

Tribe 3. *Linnaeae*. *Leaves* undivided. *Anthers* opening inwards. *Carpels* (3-4), of which two have ∞ sterile *ovules* and 1-2 have one fertile *ovule*. *Style* long.

Stems slender, creeping. Peduncles 2-flowered. Ovary 3-loc. **Linnaea.**

Shrubs. Flowers in racemes. Ovary 4-loc.

***Symphoricarpus.**

Tribe 4. *Lonicereae*. *Leaves* usually undivided. *Anthers* introrse. *Carpels* (2-5-8) all with ∞ *ovules*. *Style* usually very long.

Only British genus (Flowers zygomorphic):

Lonicera.

Fam. 3. ADOXACEAE. A small succulent herb. *Leaves* ternate. *Flowers* in 5-flowered heads, terminal 4, lateral 5-merous, homochlamydeous. *Calyx* (*bracts and bracteoles*) 2-3-lobed. *Stamens* 4 or 5-6, split to the base. *Carpels* (3-5), each with one pendulous *ovule*. *Fruit* a drupe with 1-3 *stones*. The position of this monotypic genus is very uncertain.

Only genus:

Adoxa.

B. *Stamens* fewer than *corolla-lobes*. *Ovary* inferior always with only one fertile *loculus* with one pendulous *ovule*. (For A see page 99.)

Fam. 4. VALERIANACEAE. Herbs, rarely shrubs, with opposite exstipulate *leaves*. *Inflorescence* cymose. *Flowers* rather irregular. *Calyx* usually an involute pappus expanding as fruit ripens. *Corolla* (5) or (3-4), often with pouch or spur. *Stamens* 1-4. *Anthers* exserted, versatile. *Carpels* (3), of which one has one pendulous anatropous *ovule*, the other two empty. *Style* 1. *Stigmas* 1-3.

Tribe *Valerianeae*. *Stamens* usually 3, more rarely 2 or 1.

Calyx obscurely 5-toothed. *Stamens* 3.

Valerianella.

Calyx forming pappus. *Corolla-tube* pouched at base. *Stamens* 3.

Valeriana.

Calyx forming pappus. *Corolla-tube* spurred at base. *Stamen* 1.

***Kentranthus.**

Fam. 5. DIPSACACEAE. Usually herbs. *Leaves* opposite, exstipulate. *Flowers* usually in heads. *Outer bracts* often forming involucre, *inner* axillant to flowers or absent. *Bracteoles* forming *epicalyx* (*involucel*). *Flowers* hermaphrodite, mostly zygomorphic. *Stamens* 4 or fewer. *Carpels* (2), but *ovary* 1-loc with one pendulous *ovule*. *Style* 1. *Stigmas* 1-2. *Fruit* indehiscent, enclosed in *epicalyx*.

Involucral-bracts very long, rigid. *Floral-bracts* spinous, longer than flowers.

Dipsacus.

Involucral-bracts short. *Floral-bracts* short or absent.

Scabiosa.

β. *Stamens conniving or partly united.* (For α see p. 98.)

Order 8. **Cucurbitales.** Mostly climbing herbs. *Flowers* typically 5-merous. *Stamens* 5, rarely all free, usually appearing as 3 by union of two pairs, or all 5 may be united into a *synandrium*.

Fam. CUCURBITACEAE. Usually annual herbs of rapid growth climbing by means of *tendrils*. *Leaves* spiral, mostly lobed. *Flowers* cymose, usually diclinous, actinomorphic. *Flower-axis* cup-shaped. *Stamens* 5 free (rare), or more often apparently 3 by fusion. *Anthers* often twisted. *Carpels* usually (3), each with two axile revolute *placentae* bearing usually ∞ anatropous *ovules*. *Ovary* inferior, usually 3-locular. *Style* 1. *Stigmas* usually 3. *Fruit* usually a berry. No *endosperm*.

Tribe *Cucurbiteae*. *Stamens* 5, mostly two pairs united and one free. *Anthers* with ω -shaped or U-shaped *pollen-sacs*.

Only British genus:

Bryonia.

Order 9. **Campanulatae.** Mostly herbs. *Flowers* typically 5-merous with isomerous *androecium* and oligomerous *gynaecium*. *Anthers* often united. *Ovary* inferior, with several *loculi* each with $\infty-1$ *ovules*, or 1-loc with one *ovule*.

Fam. 1. CAMPANULACEAE. Usually herbs with latex. *Leaves* spiral, exstipulate. *Flowers* often large, usually 5-merous, hermaphrodite, and actinomorphic. *Stamens* free or united. *Anthers* introrse. *Carpels* mostly (2-5) with ∞ anatropous *ovules*. *Style* 1, often with device for brushing up and collecting *pollen*. *Ovary* mostly inferior, of several *loculi*, rarely 1-loc. *Placentae* usually axile. *Fruit* usually capsular. *Seeds* with fleshy *endosperm*.

Subfamily 1. *Campanuloideae*. Flowers actinomorphic, rarely slightly zygomorphic. Anthers usually free.

Tribe *Campanuleae*. Flowers symmetrical. Corolla-lobes valvate in bud.

Subtribe 1. *Campanulinae*. Ovary always quite inferior. Fruit usually a capsule dehiscing laterally.

Flowers often capitate. Corolla-lobes linear.

Phyteuma.

Flowers racemose. Corolla-lobes broad. Capsule short.

Campanula.

As *Campanula*, but ovary and capsule slender.

Specularia.

Subtribe 2. *Wahlenbergiinae*. Ovary inferior or almost superior. Fruit usually a capsule dehiscing apically.

Flowers capitate. Corolla-lobes almost free. Anthers united at base.

Jasione.

Inflorescence lax. Corolla campanulate. Anthers free.

Wahlenbergia.

Subfamily 2. *Lobelioideae*. Flowers zygomorphic and resupinate. Anthers united.

Only British genus :

Lobelia.

Fam. 2. COMPOSITAE. Mostly herbs. Leaves spiral or opposite, exstipulate, sometimes with auricles. Flowers in involucrate heads. Involucral-bracts spiral or whorled, sometimes united, usually not bearing flowers in their axils. Floral-bracts chaffy, setaceous, petaloid, or absent. Flowers 5-merous, hermaphrodite, or ♂ and ♀ flowers

separate, or sterile, actinomorphic or zygomorphic. *Calyx* rarely distinctly developed, usually represented by hairs, bristles or scales forming the *pappus*. *Corolla* tubular, ligulate, or 2-lipped (in Tribe *Mutisieae*, not represented in Britain). *Stamens* inserted on the *corolla-tube*, *filaments* usually free. *Anthers* usually united, forming a tube, dehiscing inwards. *Carpels* (2), median, but the *ovary* is 1-locular with one ascending, anatropous *ovule*. *Style* bifid in the fertile flowers, the two arms stigmatic on their inner surfaces. *Fruit* a 1-seeded, inferior nut. *Seeds* without endosperm.

Subfamily 1. *Tubuliflorae*. No *latex* present. *Corolla* of *disc-flowers* never ligulate.

Tribe 1. *Eupatorieae*. *Leaves* usually opposite. *Flowers* all hermaphrodite and alike. *Corolla* tubular, never pure yellow. *Anthers* basifixed, obtuse at base. *Style-arms* hairy outside, but the style has no ring of hair below the bifurcation.

Only British genus :

Eupatorium.

Tribe 2. *Astereae*. *Leaves* spiral. *Heads* mostly with ♀ ligulate *ray-flowers* and ♂ tubular *disc-flowers*. *Anthers* as in *Eupatorieae*. *Style* without ring of hairs below bifurcation, arms thickly hairy above.

Ray-flowers 1-seriate, yellow: disc-flowers yellow. *Pappus* scabrid.

Solidago.

Ray-flowers 1-seriate, white or pink: disc-flowers yellow. *Pappus* absent.

Bellis.

Ray-flowers 1-seriate, white, blue or purple: disc-flowers yellow. *Pappus* many-seriate, persistent. **Aster.**

As *Aster*, but ray-flowers 2 or more seriate.

Erigeron.

Tribe 3. *Inuleae*. *Leaves* spiral. *Ray-flowers* mostly ♀: *disc-flowers* often 4-merous. *Anthers* with tails at the base. *Pappus* mostly of hairs.

1. Corolla of ♀ (ray) flowers filiform. Chaffy floral-bracts present. **Filago.**

2. As 1 but floral-bracts absent.

Heads dioecious.

Antennaria (including ***Anaphalis**).

Heads with ♂ and ♀ flowers. **Gnaphalium.**

3. Corolla of ♀ (ray) flowers ligulate. Floral-bracts absent.

Pappus of scabrid hairs. **Inula.**

Pappus of an inner row of scabrid hairs and an outer row of scales. **Pulicaria.**

Tribe 4. *Heliantheae*. *Leaves* usually opposite. *Involucral-bracts* without broad scarious margin. *Disc-flowers* actinomorphic. *Floral-bracts* present. *Anthers* basifixed with obtuse base. *Style* with hairs mostly above bifurcation. *Pappus* never of hairs.

Subtribe 1. *Ambrosiinae*. Corolla of ♀ flowers reduced or absent. *Anthers* sometimes quite free.

Xanthium.

Subtribe 2. *Coreopsidinae*. All flowers with perfect corollas. Corolla of ray-flowers early deciduous. *Pappus* absent or of bristles. **Bidens.**

Subtribe 3. *Galinsoginae*. As above, but *pappus* of ciliate scales. ***Galinsoga.**

Tribe 5. *Anthemideae*. As *Heliantheae* but often aromatic, *leaves* spiral, and *involucral-bracts* with broad scarious margins. *Pappus* minute or absent.

Subtribe 1. *Anthemidinae*. Floral-bracts present.

Ray-flowers with oblong ligules. Fruit not much compressed. **Anthemis.**

Ray-flowers with short, broad ligule. Fruit much compressed. **Achillea.**

Ray-flowers absent. Corolla compressed, with corky appendages. **Diotis.**

Subtribe 2. *Chrysantheminae*. Floral-bracts absent.

Ray-flowers ligulate. Involucral-bracts 1-2-seriate, all equal. **Matricaria.**

Ray-flowers ligulate. Involucral-bracts many-seriate, outer shorter. **Chrysanthemum.**

Flowers all tubular, 4-merous. ***Cotula.**

Flowers all tubular, 5-merous. Axis of head broad. **Tanacetum.**

Flowers all tubular, 5-merous. Axis of head narrow. **Artemisia.**

Tribe 6. *Senecioneae*. Leaves spiral. Involucral-bracts usually 1-seriate, without scarious margin. Flowers usually all yellow. Floral-bracts absent. Anthers usually obtuse at base. Style-arms various. Pappus usually of copious, soft hairs.

* Leaves often very large, produced after the flowers. Style-arms of disc-flowers connate.

Heads in racemes, purple or white, outer flowers tubular. **Petasites.**

Heads solitary, yellow, outer flowers ligulate.

Tussilago.

** Leaves radical and cauline. Style-arms of disc-flowers separate, truncate.

Perennial herbs. Involucral-bracts in several series all equal. ***Doronicum.**

Annual and perennial herbs. Involucral-bracts 1-seriate or with a few smaller ones at the base.

Senecio.

*Tribe 7. *Calenduleae*. Leaves spiral. Axis of head naked. Disc-flowers hermaphrodite, functionally ♂ (not ripening fruit). Ray-flowers ♀. Anthers pointed at base. Pappus absent.

(Fruit bent inwards.)

***Calendula.**

Tribe 8. *Cynareae*. Leaves spiral, often spinous. Involucral-bracts many-seriate, often spinous. Axis of head usually with bristles which have no definite relationship to the flowers. Flowers often purple, usually all tubular, all hermaphrodite, or outer ♀ or neuter. Anthers usually with tails at base. Style usually with ring of hairs or swelling.

Subtribe 1. *Echinopsinae*. Heads 1-flowered, themselves arranged in heads of the second order. Flowers blue or white.

***Echinops.**

Subtribe 2. *Carlininae*. Heads many-flowered. Fruit with straight insertion, usually covered with silky hairs. Pappus of scales or of 1-seriate, feathery hairs.

Only British genus:

Carlina.

Subtribe 3. *Carduinae*. Heads many-flowered. Fruit with straight insertion, usually glabrous. Pappus usually of many-seriate hairs.

A. Pappus hairs simple.

Outer involucral-bracts hooked.

Arctium.

Outer involucral-bracts not hooked.

Filaments free, warted or hairy. Pappus hairs connate at base.

Carduus.

Filaments connate into a sheath. Pappus free.

Silybum.

B. Pappus hairs all, or inner ones only, feathery.

Involucral-bracts without spines. Filaments glabrous.

Saussurea.

Involucral-bracts with spines. Filaments warty or hairy. Fruit compressed or terete.

Cnicus.

Involucral-bracts with spines. Filaments nearly glabrous. Fruit 4-angled.

Onopordon.

Subtribe 4. *Centaureinae*. Heads many-flowered. Fruit with oblique insertion, glabrous or hairy. Pappus many-seriate.

Involucral-bracts without appendages. All flowers hermaphrodite.

Serratula.

Involucral-bracts with scarious or thorny appendages. Outer flowers quite sterile and larger than inner ones.

Centaurea.

Subfamily 2. *Liguliflorae*. Latex present. Corolla of all flowers ligulate.

Tribe 9. *Cichorieae*. Leaves spiral. Style-arms cylindrical, hairy, obtuse.

Subtribe 1. *Hyoseridinae*. Fruit obtuse. Pappus of scales. (Flowers blue in British species.)

Cichorium.

Subtribe 2. *Lapsaninae*. Involucral-bracts almost equal. Fruit obtuse. Pappus absent. (Flowers yellow in British species.)

Leaves all radical. Fruit crowned with ring.

Arnoseris.

Leaves radical and cauline. Fruit without ring.

Lapsana.

Subtribe 3. *Crepidinae*. Neither stellate nor woolly hairs present. Only inner involucral-bracts equal. Fruit with simple or feathery pappus.

Pappus of feathery hairs. **Picris.**

Pappus of simple hairs. **Crepis.**

Subtribe 4. *Hieraciinae*. Stellate or woolly hairs often present on leaves. Fruit rounded at apex. Pappus rough, brown. **Hieracium.**

Subtribe 5. *Hypochaeridinae*. Involucral-bracts all imbricate, the inner ones keeled after flowering. Fruit beaked. Pappus simple or feathery.

Narrow, membranous floral-bracts present. Beak of inner fruits long. Pappus hairs feathery. **Hypochaeris.**

No floral-bracts. All beaks short. Pappus hairs feathery. **Leontodon.**

No floral-bracts. All beaks long. Pappus hairs simple. **Taraxacum.**

Subtribe 6. *Lactucinae*. Leaves usually pinnatifid. Inner involucral-bracts equal and unaltered after flowering. Fruit acute or beaked, with numerous simple pappus hairs.

Fruit not beaked, scarcely narrowed above. **Sonchus.**

Fruit narrowed or beaked above. **Lactuca.**

Subtribe 7. *Scorzonerinae*. Leaves linear, entire. Fruit usually with long beak. Pappus simple or feathery.

Only British genus (Involucral-bracts 1-seriate):

Tragopogon.

INDEX

- Abies 9
- Abietineae, tribe 9
- Acalypheae, tribe 64
- Acer 66
- Aceraceae, family 66
- Aceras 31
- Achillea 106
- Aconitum 47
- Acoreae, tribe 22
- Acorus 22
- Actaea 47
- Adiantinae, subtribe 6
- Adiantum 6
- Adonis 48
- Adoxa 100
- Adoxaceae, family 100
- Aegopodium 78
- Aesculus 66
- Aethusa 79
- Agrimonia 59
- Agropyrum 20
- Agrostemma 44
- Agrostideae, tribe 17
- Agrostis 17
- Aira 18
- Aizoaceae, family 42
- Ajuga 92
- Ajugeae, tribe 91
- Ajugoideae, subfamily 91
- Alchemilla 59
- Alisma 13
- Alismataceae, family 13
- Alismatales (Fluviales, Helobiae),
order 11
- Alismatineae, suborder 13
- Allieae, tribe 25
- Allioideae, subfamily 25
- Allium 26
- Alnus 35
- Alopecurus 17
- Alsineae, tribe 43
- Alsinoideae, subfamily 43
- Althaea 68
- Alysseae (Latiseptae pleurorhizae),
tribe 52
- Alyssum 52
- Amarantaceae, family 41
- Amarantus 41
- Amaryllidaceae, family 27
- Amaryllideae, tribe 28
- Amaryllidoideae, subfamily 27
- Ambrosiinae, subtribe 105
- Amentiflorae (Moss) 33
- Ammineae, tribe 77
- Ammophila 17
- Anagallidinae, subtribe 84
- Anagallis 84
- Anaphalis 105
- Anchusa (including Lycopsis) 90
- Anchuseae, tribe 90
- Andromeda 82
- Andromedeae, tribe 82
- Androsaceae, tribe 83
- Anemone 48
- Anemoneae, tribe 47
- Angelica (including Archangelica)
79
- Angelicinae, subtribe 79
- Angiospermae, subdivision 10
- Angustiseptae diplecobae, tribe
53
- Angustiseptae pleurorhizae, tribe
52
- Antennaria (including Anaphalis)
105
- Anthemideae, tribe 105
- Anthemidinae, subtribe 106
- Anthemis 106
- Anthoxanthum 17
- Anthriscus 77

- Anthyllis 61
 Antirrhineae, tribe 95
 Antirrhinoideae, subfamily 95
 Antirrhinum 96
 Apera 17
 Apioideae, subfamily 76
 Apium 78
 Apocynaceae, family 87
 Aponogeton 12
 Aponogetonaceae, family 12
 Aquifoliaceae, family 65
 Aquilegia 47
 Arabideae (Siliquosae pleurorhizae),
 tribe 50
 Arabis 51
 Araceae, family 22
 Araliaceae, family 74
 Arbuteae, tribe 82
 Arbutoideae, subfamily 82
 Arbutus 82
 Archangelica 79
 Archegoniatae, division 1
 Archichlamydeae, subclass 33
 Arctium 107
 Arctostaphylos 82
 Areae, tribe 22
 Arenaria 43
 Aristolochia 39
 Aristolochiaceae, family 38
 Aristolochiales, order 38
 Aristolochieae, tribe 39
 Armeria 85
 Arnoseris 108
 Aroideae, subfamily 22
 Arrhenatherum 18
 Artemisia 106
 Arum 22
 Asareae, tribe 39
 Asarum 39
 Asparageae, tribe 27
 Asparagoideae, subfamily 26
 Asparagus 27
 Asperugo 90
 Asperula 99
 Asphodeloideae, subfamily 25
 Aspidieae, tribe 6
 Asplenieae, tribe 6
 Aspleninae, subtribe 6
 Asplenium 6
 Aster 104
 Astereae, tribe 104
 Astragaleae, tribe 61
 Astragalus 61
 Astrantia 76
 Athyrium 6
 Atriplex 41
 Atripliceae, tribe 41
 Atropa 94
 Avena 18
 Aveneae, tribe 18
 Azalea 82
 Azolla 7
 Ballota 93
 Balsaminaceae, family 67
 Balsaminineae, suborder 66
 Barbarea 51
 Bartschia 97
 Bellis 104
 Berberidaceae, family 48
 Berberideae, tribe 48
 Berberidoideae, subfamily 48
 Berberis 48
 Beta 41
 Beteae, tribe 41
 Betula 35
 Betulaceae, family 35
 Betuleae, tribe 35
 Bidens 105
 Blackstonia (= Chlora) 86
 Blechninae, subtribe 6
 Blechnum 6
 Borraginaceae, family 89
 Borraginineae, suborder 89
 Borraginoideae, subfamily 89
 Borrigo 90
 Botrychium 8
 Brachycarpeae (Angustiseptae di-
 plecolobae), tribe 53
 Brachypodium 19
 Brassica 51
 Brassiceae (Siliquosae orthoplocae),
 tribe 51
 Briza 19
 Bromus 19
 Brunella 93
 Brunellinae, subtribe 93
 Bryonia 102
 Bryophyta, subdivision 1
 Bupleurum 78
 Butomaceae, family 14
 Butomineae, suborder 14
 Butomus 14
 Buxaceae, family 65

- Buxineae, suborder 65
 Buxus 65
 Cakile 53
 Cakilinae (Lomentaceae pleuro-
 rhizae), tribe 53
 Calamagrostis 17
 Calamintha 94
 Calendula 107
 Calenduleae, tribe 107
 Callitrichaceae, family 64
 Callitriche 64
 Callitrichinae, suborder 64
 Calluna 83
 Caltha 47
 Calystegia 88
 Camelina 52
 Camelineae (Latiseptae notorhizae),
 tribe 52
 Campanula 103
 Campanulaceae, family 102
 Campanulatae, order 102
 Campanuleae, tribe 103
 Campanulinae, subtribe 103
 Campanuloideae, subfamily 103
 Cannabaceae, family 36
 Cannabis 37
 Capparidinae, suborder 50
 Caprifoliaceae, family 99
 Capsella 53
 Cardamine 51
 Carduinae, subtribe 107
 Carduus 107
 Carex 21
 Caricoideae, subfamily 21
 Carinae, subtribe 77
 Carlina 107
 Carlininae, subtribe 107
 Carpinus 35
 Carum 78
 Caryophyllaceae, family 43
 Caryophyllinae, suborder 43
 Castanea 36
 Castaneae, tribe 36
 Catabrosa 19
 Caucalinae, subtribe 77
 Caucalis (including Torilis) 77
 Cedrus 10
 Celastraceae, family 65
 Celastrineae, suborder 65
 Centaurea 108
 Centaureinae, subtribe 108
 Centaurium 87
 Centrospermae, order 40
 Centunculus 84
 Cephalanthera 32
 Cephalantherinae, subtribe 32
 Cerastium (including Moenchia) 43
 Ceratophyllaceae, family 46
 Ceratophyllum 46
 Chaerophyllum 77
 Cheilanthinae, subtribe 6
 Cheiranthus 51
 Chelidoniae, tribe 49
 Chelidonium 49
 Cheloneae, tribe 96
 Chenopodiaceae, family 40
 Chenopodieae, tribe 41
 Chenopodinae, suborder 40
 Chenopodium 41
 Chironieae, tribe 86
 Chlora 86
 Chlorideae, tribe 19
 Chrysantheminae, subtribe 106
 Chrysanthemum 106
 Chrysosplenium 56
 Cicendia 87
 Cichorieae, tribe 108
 Cichorium 108
 Cicuta 78
 Circaea 73
 Cistaceae, family 71
 Cistineae, suborder 70
 Cladium 21
 Claytonia 42
 Clematis 47
 Cnicus 108
 Cochlearia 52
 Coffeoidaeae, subfamily 99
 Colchiceae, tribe 25
 Colchicum 25
 Compositae, family 103
 Coniferae, class 9
 Conium 77
 Conopodium 76
 Contortae, order 85
 Convallaria 27
 Convallarieae, tribe 27
 Convallariinae, subtribe 27
 Convolvulaceae, family 88
 Convolvuleae, tribe 88
 Convolvulinae, suborder 88
 Convolvuloideae, subfamily 88
 Convolvulus 88

- Corallorhiza 32
 Coreopsidinae, subtribe 105
 Coriandreae, tribe 77
 Coriandrum 77
 Cornaceae, family 80
 Cornoideae, subfamily 80
 Cornus 80
 Coronilla 61
 Coronopus 53
 Corrigiola 44
 Corydalis 50
 Coryleae, tribe 35
 Corylus 35
 Corynephorus 18
 Cotoneaster 58
 Cotula 106
 Cotyledon 55
 Crambe 53
 Crassula (including Tillaea) 55
 Crassulaceae, family 55
 Crataegus 58
 Crepidinae, subtribe 109
 Crepis 109
 Crithmum 78
 Crocoideae, subfamily 29
 Crocus 29
 Crotonoideae, subfamily 64
 Cruciferae, family 50
 Cryptogramme 6
 Cucubalus 44
 Cucurbitaceae, family 102
 Cucurbitales, order 102
 Cucurbiteae, tribe 102
 Cupressineae, tribe 10
 Cuscuta 88
 Cuscutoideae, subfamily 88
 Cyclamen 84
 Cyclamineae, tribe 84
 Cyclolobeae (Chenopodiaceae),
 family 41
 Cymbalaria 96
 Cynareae, tribe 107
 Cynodon 20
 Cynoglosseae, tribe 89
 Cynoglossum 89
 Cynosurus 18
 Cyperaceae, family 20
 Cyperinae, subtribe 21
 Cyperus 21
 Cypripedileae, tribe 31
 Cypripedium 31
 Cystopteris 5
 Cytisus 60
 Dabeocia 82
 Dactylis 19
 Damasonium 13
 Daphne 72
 Datura 95
 Datureae, tribe 95
 Dauceae, tribe 79
 Daucus 79
 Delphinium 47
 Dentaria 51
 Deschampsia 18
 Deyeuxia 17
 Diandrae, subfamily 30
 Diantheae, tribe 45
 Dianthus (including Tunica) 45
 Dicotyledones, class 33
 Digitaleae, tribe 96
 Digitalis 96
 Dioscoreaceae, family 28
 Dioscoreae, tribe 28
 Diotis 106
 Diplotaxis 52
 Diplozygieae (Umbelliferae), family
 79
 Dipsacaceae, family 101
 Dipsacus 101
 Doronicum 106
 Draba (including Erophila) 52
 Drosera 54
 Droseraceae, family 54
 Dryas 58
 Dryopteris (including Phegopteris) 6
 Echieae, tribe 90
 Echinodorus 13
 Echinops 107
 Echinopsinae, subtribe 107
 Echium 90
 Elaeagnaceae, family 72
 Elatinaceae, family 70
 Elatine 70
 Elatinoides 96
 Eligulatae, suborder 2
 Elisma 13
 Elodea 15
 Elymus 20
 Embryophyta Asiphonogama
 (Archegoniatae), division 1
 Embryophyta Siphonogama,
 division 8

- Empetraceae, family 65
 Empetrineae, suborder 65
 Empetrum 65
 Epilobiaceae (Onagraceae), family 73
 Epilobium 73
 Epimediaceae, tribe 48
 Epimedium 48
 Epipactis 32
 Epipogon 32
 Equisetaceae, family 2
 Equisetales, order 2
 Equisetariae, class 1
 Equisetum 2
 Eranthis 47
 Erica 83
 Ericaceae, family 81
 Ericales, order 80
 Ericaceae, tribe 83
 Ericoideae, subfamily 83
 Erigeron 104
 Erinus 96
 Eriocaulaceae, family 23
 Eriocauloideae, subfamily 24
 Eriocaulon 24
 Eriophorum 21
 Eritrichiaceae, tribe 89
 Erodium 62
 Erophila 52
 Eryngium 76
 Erysimum 51
 Eulineae, tribe 63
 Euonymus 66
 Eupatorieae, tribe 104
 Eupatorium 104
 Euphorbia 64
 Euphorbiaceae, family 63
 Euphorbieae, tribe 64
 Euphrasia 97
 Fagaceae, family 35
 Fagales, order 34
 Fageae, tribe 36
 Fagopyrum 40
 Fagus 36
 Farinosae, order 23
 Ferulinae, subtribe 79
 Festuca 19
 Festuceae, tribe 18
 Filago 105
 Filicales, order 4
 Filicariae, class 3
 Filicineae, suborder 4
 Flacourtiaceae, suborder 71
 Fluviales, order 11
 Foeniculum 78
 Fragaria 58
 Frankenia 70
 Frankeniaceae, family 70
 Fraxineae, tribe 86
 Fraxinus 86
 Fritillaria 26
 Fuchsia 73
 Fumaria 50
 Fumarioideae, subfamily 50
 Gagea 25
 Galanthinae, subtribe 28
 Galanthus 28
 Galeopsis 93
 Galieae, tribe 99
 Galinsoga 105
 Galinsoginae, subtribe 105
 Galium 99
 Gastridium 17
 Genista 60
 Genisteae, tribe 60
 Gentiana 87
 Gentianaceae, family 86
 Gentianineae, suborder 86
 Gentianoideae, subfamily 86
 Geraniaceae, family 62
 Geraniales, order 61
 Geranieae, tribe 62
 Geraniineae, suborder 62
 Geranium 62
 Geum 59
 Gladioleae, tribe 29
 Gladiolus 29
 Glaucium 49
 Glaux 84
 Glumiflorae, order 15
 Glyceria 19
 Gnaphalium 105
 Goodyera 32
 Gradatae (Bower) 5
 Graminaceae, family 15
 Gratiroleae, tribe 96
 Guttiferae, family 69
 Gymnadeniinae, subtribe 31
 Gymnogramme 6
 Gymnogramminae, subtribe 6
 Gymnospermae, subdivision 8
 Gynandreae, suborder 30

- Habenaria 31
 Halorrhagaceae, family 74
 Halorrhagoideae, subfamily 74
 Haplostemonous (Crassulaceae), family 55
 Haplozygiaeae (Umbelliferae), family 76
 Hedera 75
 Hedysareae, tribe 61
 Heleocharis 21
 Heliantheae, tribe 105
 Helianthemum 71
 Helleboreae, tribe 47
 Helleborus 47
 Helobiae, order 11
 Heracleum 79
 Herminium 31
 Herniaria 44
 Hesperis 51
 Heterochlamydeae (Moss) 45
 Hieraciinae, subtribe 109
 Hieracium 109
 Hierochloë 17
 Hippocastanaceae, family 66
 Hippocrepis 61
 Hippophaë 72
 Hippuridaceae, family 74
 Hippuridineae, suborder 74
 Hippuris 74
 Holcus 18
 Holosteam 43
 Hordeae, tribe 20
 Hordeum 20
 Hottonia 84
 Hottoniinae, subtribe 84
 Humulus 37
 Hutchinsia 53
 Hyacinthus 26
 Hydrilleae, tribe 15
 Hydrocharis 15
 Hydrocharitaceae, family 14
 Hydrochariteae, tribe 15
 Hydrocotyle 75
 Hydrocotyleae, tribe 75
 Hydrocotyloideae, subfamily 75
 Hydropteridineae, suborder 7
 Hymenophyllaceae, family 5
 Hymenophyllum 5
 Hyoscyamus 95
 Hyoseridinae, subtribe 108
 Hypericeae, tribe 69
 Hypericoideae, subfamily 69
 Hypericum 69
 Hypochaeridinae, subtribe 109
 Hypochaeris 109
 Iberis 52
 Ilex 65
 Illecebrum 44
 Impatiens 67
 Inula 105
 Inuleae, tribe 105
 Iridaceae, family 28
 Iridineae, suborder 28
 Iridoideae, subfamily 29
 Iris 29
 Isatideae (Nucamentaceae notorhizae), tribe 53
 Isatis 53
 Isoëtaceae, family 3
 Isoëtales, order 3
 Isoëtes 3
 Ixioideae, subfamily 29
 Jasione 103
 Juglandaceae, family 34
 Juglandales, order 34
 Juglans 34
 Juncaceae, family 24
 Juncaginaceae, family 13
 Juncineae, suborder 24
 Juncus 24
 Juniperus 10
 Kentranthus 101
 Kobresia 21
 Koeleria 19
 Labiatae, family 91
 Lactuca 109
 Lactucinae, subtribe 109
 Lagurus 17
 Lamiinae, subtribe 93
 Lamium 93
 Lappula 90
 Lapsana 108
 Lapsaninae, subtribe 108
 Larix 10
 Lathraea 97
 Lathyrus 61
 Latiseptae diplecobolae, tribe 52
 Latiseptae notorhizae, tribe 52
 Latiseptae pleurorhizae, tribe 52
 Lavatera 68

- Ledeae, tribe 82
 Ledum 82
 Leersia 16
 Leguminosae, family 59
 Lemna 23
 Lemnaceae, family 23
 Lemnoideae, subfamily 23
 Lentibulariaceae, family 97
 Leontodon 109
 Leonurus 93
 Lepidieae, tribe 53
 Lepidium 53
 Lepturus 20
 Leucojum 28
 Ligulatae, suborder 3
 Liguliflorae, subfamily 108
 Ligusticum 79
 Ligustrum 86
 Liliaceae, family 24
 Liliiflorae, order 24
 Liliineae, suborder 24
 Lilioideae, subfamily 26
 Lilium 26
 Limosella 96
 Linaceae, family 63
 Linaria 96
 Linnaea 100
 Linnaeae, tribe 100
 Linum 63
 Liparideae, tribe 32
 Liparis 32
 Listera 32
 Lithospermeae, tribe 90
 Lithospermum 90
 Litorella 98
 Lloydia 26
 Lobelia 103
 Lobelioideae, subfamily 103
 Loiseluria 82
 Lolium 20
 Lomentaceae orthoplocae, tribe 53
 Lomentaceae pleurorhizae, tribe 53
 Lonicera 100
 Lonicereae, tribe 100
 Loranthaceae, family 38
 Lorantheae, suborder 38
 Loteae, tribe 60
 Lotus 61
 Ludwigia 73
 Lupinus 60
 Luzula 24
 Lychnideae, tribe 44
 Lychnis 44
 Lycium 94
 Lycopodiaceae, family 2
 Lycopodiales, order 2
 Lycopodiariae, class 2
 Lycopodium 2
 Lycopsis 90
 Lycopus 94
 Lysimachia 84
 Lysimachieae, tribe 84
 Lysimachiinae, subtribe 84
 Lythraceae, family 72
 Lythrae, tribe 73
 Lythrum 73
 Maianthemum 27
 Malaxis 32
 Malva 68
 Malvaceae, family 68
 Malvales, order 67
 Malveae, tribe 68
 Malvineae, suborder 68
 Marrubieae, tribe 92
 Marrubium 92
 Marsiliaceae, family 7
 Matricaria 106
 Matthiola 51
 Meconopsis 49
 Medicago 60
 Melampyrum 97
 Melandryum 44
 Melanthioideae, subfamily 25
 Melica 19
 Melilotus 60
 Melissa 94
 Melittinae, subtribe 93
 Melittis 93
 Mentha 94
 Menyanthes 87
 Menyanthoideae, subfamily 87
 Mercurialis 64
 Mertensia 90
 Mesembryanthemeae, tribe 42
 Mesembryanthemum 42
 Mespilus 58
 Metachlamydeae (Sympetalae), sub-
 class 80
 Meum 79
 Mibora 17
 Microcala 87
 Microspermae, order 29
 Milium 17

- Mimulus 96
 Mixtae (Bower) 5
 Moenchia 43
 Molinia 19
 Monandreae, subfamily 31
 Moneses 81
 Monocotyledones, class 10
 Monotropa 81
 Monotropeae, tribe 81
 Monotropeoideae, subfamily 81
 Montia 42
 Moraceae, tribe 29
 Muscari 26
 Myosotis 90
 Myosurus 48
 Myrica 34
 Myricaceae, family 34
 Myricales, order 34
 Myriophyllum 74
 Myrrhis 77
 Myrtiflorae, order 71
 Myrtineae, suborder 72

 Naiadaceae, family 12
 Naias 12
 Narcisseae, tribe 28
 Narcissinae, subtribe 28
 Narcissus 28
 Nardus 20
 Narthecium 25
 Nasturtium 51
 Neottia 32
 Neottieae, tribe 32
 Nepeta 92
 Nepeteae, tribe 92
 Nucamentaceae notorhizae, tribe 53
 Nuphar 46
 Nuphareae, tribe 46
 Nymphaea 46
 Nymphaeaceae, family 45
 Nymphaeaceae, suborder 45
 Nymphaeaoideae, subfamily 45

 Obdiplostemonous (Crassulaceae),
 family 55
 Oenanthe 79
 Oenothera 73
 Oenotheraeae, tribe 73
 Oleaceae, family 85
 Oleaceae, tribe 86
 Oleineae, suborder 85
 Oleoideae, subfamily 86

 Omphalodes 89
 Onagraceae, family 73
 Onobrychis 61
 Ononis 60
 Onopordon 108
 Ophioglossaceae, family 8
 Ophioglossales, order 7
 Ophioglossum 8
 Ophrydeae, tribe 31
 Ophrys 31
 Orchidaceae, family 30
 Orchis 31
 Origanum 94
 Ornithogalum 26
 Ornithopus 61
 Orobanchaceae, family 97
 Orobanche 97
 Oryzeae, tribe 16
 Osmunda 5
 Osmundaceae, family 4
 Oxalidaceae, family 63
 Oxalis 63
 Oxycoccus 82
 Oxyria 39
 Oxytropis 61

 Paeonia 46
 Paeonieae, tribe 46
 Pandanales, order 10
 Paniceae, tribe 16
 Panicoideae, subfamily 16
 Panicum 16
 Papaver 49
 Papaveraceae, family 49
 Papavereae, tribe 49
 Papaveroideae, subfamily 49
 Papilionatae, subfamily 59
 Parideae, tribe 27
 Parietales, order 69
 Parietaria 37
 Parietarieae, tribe 37
 Paris 27
 Parnassia 56
 Parnassieae, tribe 56
 Paronychieae, tribe 44
 Pedicularis 97
 Peplis 73
 Petaloideae (Moss) 37
 Petasites 106
 Peucedaneae, tribe 79
 Peucedanum 79
 Phalarideae, tribe 16

- Phalaris* 16
Phanerogamae, division 8
Phegopteris 6
Phleum 17
Phragmites 18
Phyllodoce 82
Phyllodoceae, tribe 82
Physospermum 77
Physurinae, subtribe 32
Phyteuma 103
Phytolaccineae, suborder 42
Picea 9
Picris 109
Pilularia 7
Pimpinella 78
Pinaceae, family 9
Pinguicula 98
Pinus 10
Pirola (including *Moneses*) 81
Pirolaceae, family 80
Pirolloideae, subfamily 81
Pirus 58
Plantaginaceae, family 98
Plantaginales, order 98
Plantago 98
Platanaceae, family 56
Platanus 57
Pleonandrae (*Diandrae*), subfamily 30
Plumbaginaceae, family 85
Plumbaginales, order 84
Plumerieae, tribe 87
Plumerioideae, subfamily 87
Poa 19
Poëoideae, subfamily 16
Polemoniaceae, family 88
Polemonieae, tribe 89
Polemonioidae, subfamily 89
Polemonium 89
Polycarpeae, tribe 43
Polycarpon 43
Polygala 63
Polygalaceae, family 63
Polygalineae, suborder 63
Polygonaceae, family 39
Polygonales, order 39
Polygonateae, tribe 27
Polygonatum 27
Polygoneae, tribe 40
Polygonoideae, subfamily 39
Polygonum (including *Fagopyrum*) 40
Polypodiaceae, family 5
Polypodieae, tribe 7
Polypodium 7
Polypogon 17
Polystichum 6
Pomoideae, subfamily 58
Populus 34
Portulaca 42
Portulacaceae, family 42
Portulacineae, suborder 42
Potamogeton 12
Potamogetonaceae, family 11
Potamogetoneae, tribe 12
Potamogetonineae, suborder 11
Potentilla 58
Potentilleae, tribe 58
Poterium 59
Pothoideae, subfamily 22
Primula 83
Primulaceae, family 83
Primulales, order 83
Primulinae, subtribe 83
Prunoideae, subfamily 59
Prunus 59
Pseudosolanoideae, subfamily 95
Pseudotsuga 9
Pterideae, tribe 6
Pteridinae, subtribe 7
Pteridium (= *Pteris*) 7
Pteridophyta, subdivision 1
Pteris 7
Pulicaria 105
Pulmonaria 90
Quercus 36
Radiola 63
Ranales, order 45
Ranunculaceae, family 46
Ranunculineae, suborder 46
Ranunculus 48
Raphaneae (*Lomentaceae* orthoplocae), tribe 53
Raphanus 53
Reseda 54
Resedaceae, family 54
Resedineae, suborder 53
Rhamnaceae, family 67
Rhamnales, order 67
Rhamneae, tribe 67
Rhamnus 67
Rheum (including *Oxyria*) 39

- Rhinanthaeae, tribe 97
 Rhinanthoideae, subfamily 96
 Rhinanthus 97
 Rhododendreae, tribe 82
 Rhododendroideae, subfamily 81
 Rhododendron (including *Azalea*) 82
 Rhoecadales, order 49
 Rhoecadineae, suborder 49
 Rhynchospora 21
 Rhynchosporaeae, tribe 21
 Rhynchosporoideae, subfamily 21
 Ribes 56
 Ribesoideae, subfamily 56
 Roemeria 49
 Romulea 29
 Rosa 59
 Rosaceae, family 57
 Rosales, order 54
 Roseae, tribe 59
 Rosineae, suborder 56
 Rosoideae, subfamily 58
 Rubia 99
 Rubiaceae, family 99
 Rubiales, order 99
 Rubus 58
 Rumex 39
 Rumiceae, tribe 39
 Rumicoideae, subfamily 39
 Ruppia 12
 Ruscus 27

 Sagina 43
 Sagittaria 14
 Salicaceae, family 33
 Salicales, order 33
 Salicornia 41
 Salicornieae, tribe 41
 Salix 34
 Salsola 41
 Salsoleae, tribe 41
 Salvia 93
 Salviaeae, tribe 93
 Salviniaceae, family 7
 Sambuceae, tribe 100
 Sambucus 100
 Samoleae, tribe 84
 Samolus 84
 Sanguisorbeae, tribe 59
 Sanicula 76
 Saniculeae, tribe 76
 Saniculoideae, subfamily 76

 Santalaceae, family 38
 Santalales, order 37
 Santalineae, suborder 37
 Sapindales, order 64
 Sapindineae, suborder 66
 Saponaria 45
 Sarraceniales, order 54
 Satureia (*Calamintha*) 94
 Satureieae, tribe 93
 Saussurea 108
 Saxifraga 56
 Saxifragaceae, family 55
 Saxifrageae, tribe 56
 Saxifragineae, suborder 55
 Saxifragoideae, subfamily 56
 Scabiosa 101
 Scandicinae, subtribe 76
 Scandicineae, tribe 76
 Scandix 76
 Schefflereae, tribe 75
 Scheuchzeria 13
 Scheuchzeriaceae (*Juncaginaceae*), family 13
 Schoenus 21
 Scilla 26
 Scilleae, tribe 26
 Scirpeae, tribe 20
 Scirpinae, subtribe 21
 Scirpoideae, subfamily 20
 Scirpus 21
 Scleranthaeae, tribe 44
 Scleranthus 44
 Scolopendrium 6
 Scorzonerinae, subtribe 109
 Scrophularia 96
 Scrophulariaceae, family 95
 Scutellaria 92
 Scutellarioideae, subfamily 92
 Secale 20
 Sedum 55
 Selaginella 3
 Selaginellaceae, family 3
 Selinum 79
 Sempervivum 55
 Senecio 107
 Senecioneae, tribe 106
 Serapiadinae, subtribe 31
 Serratula 108
 Seseli 78
 Seselinae, subtribe 78
 Sesleria 18
 Setaria 16

- Sherardia 99
 Sibthorpia 96
 Sieglingia 18
 Silaus 79
 Silene 44
 Silenoideae, subfamily 44
 Siliquosae notorhizae, tribe 51
 Siliquosae orthoplocae, tribe 51
 Siliquosae pleurorhizae, tribe 50
 Silybum 108
 Simethis 25
 Simplices (Bower) 4
 Sison 78
 Sisymbrieae (Siliquosae notorhizae),
 tribe 51
 Sisymbrium 51
 Sisyrinchieae, tribe 29
 Sisyrinchium 29
 Sium 78
 Smyrnieae, tribe 77
 Smyrnum 77
 Solanaceae, family 94
 Solaneae, tribe 94
 Solanineae, suborder 94
 Solanum 95
 Solidago 104
 Sonchus 109
 Sparganiaceae, family 11
 Sparganium 11
 Spartina 20
 Spathiflorae, order 22
 Specularia 103
 Spergula 43
 Spergularia 43
 Sperguleae, tribe 43
 Spiraea 57
 Spiraeae, tribe 57
 Spiraeoideae, subfamily 57
 Spiranthes 32
 Spiranthinae, subtribe 32
 Spirodela 23
 Spirolobeae (Chenopodiaceae),
 family 41
 Stachydoideae, subfamily 92
 Stachyeae, tribe 92
 Stachys 93
 Staphylea 66
 Staphyleaceae, family 66
 Statice 85
 Staticeae, tribe 85
 Stellaria 43
 Stratioteae, tribe 14
 Stratiotes 14
 Stratiotoideae, subfamily 14
 Suaeda 41
 Suaedeae, tribe 41
 Subularia 52
 Subularieae (Latisepetae diploco-
 lobae), tribe 52
 Swertieae, tribe 87
 Sympetalae, subclass 80
 Symphoricarpus 100
 Symphytum 90
 Syringa 86
 Syringeae, tribe 86
 Tamaricaceae, family 70
 Tamariceae, tribe 70
 Tamaricineae, suborder 70
 Tamarix 70
 Tamus 28
 Tanacetum 106
 Taraxacum 109
 Taxaceae, family 9
 Taxeae, tribe 9
 Taxoideae, subfamily 9
 Taxus 9
 Teesdalia 53
 Tetrasepaleae, tribe 46
 Teucrium 92
 Thalictrum 47
 Theineae, suborder 69
 Thesieae, tribe 38
 Thesium 38
 Thlaspeae (Angustiseptae pleuro-
 rhizae), tribe 52
 Thlaspi 52
 Thymelaeaceae, family 72
 Thymelaeineae, suborder 71
 Thymelaeoideae, subfamily 72
 Thymus 94
 Tilia 68
 Tiliaceae, family 68
 Tillaea 55
 Tofieldia 25
 Tofieldieae, tribe 25
 Tordyliinae, subtribe 79
 Tordylium 79
 Torilis 77
 Tragopogon 109
 Trichomanes 5
 Tricoccae, suborder 63
 Trientalis 84
 Trifolieae, tribe 60

- Trifolium 60
 Triglochin 13
 Triglochineae, tribe 13
 Trigonella 60
 Trinia 78
 Trisetum 18
 Triticum (including Agropyrum and Secale) 20
 Trollius 47
 Tsuga (including Pseudotsuga) 9
 Tubiflorae, order 88
 Tubuliflorae, subfamily 104
 Tulipa 26
 Tulipeae, tribe 26
 Tunica 45
 Tussilago 106
 Typha 11
 Typhaceae, family 11

 Ulex 60
 Ulmaceae, family 36
 Ulmaria 59
 Ulmarieae, tribe 59
 Ulmoideae, subfamily 36
 Ulmus 36
 Umbellales (Umbelliflorae), order 74
 Umbelliferae, family 75
 Umbelliflorae, order 74
 Urereae, tribe 37
 Urtica 37
 Urticaceae, family 37
 Urticales, order 36
 Utricularia 98
 Utricularieae, tribe 98

 Vaccinieae, tribe 82
 Vaccinioideae, subfamily 82
 Vaccinium (including Oxycoccus) 82

 Valeriana 101
 Valerianaceae, family 101
 Valerianeae, tribe 101
 Valerianella 101
 Vallisnerioideae, subfamily 15
 Verbasceae, tribe 95
 Verbascum 95
 Verbena 91
 Verbenaceae, family 91
 Verbeneae, tribe 91
 Verbenineae, suborder 90
 Veronica 96
 Viburneae, tribe 100
 Viburnum 100
 Vicia 61
 Vicieae, tribe 61
 Villarsia 87
 Vinca 87
 Viola 71
 Violaceae, family 71
 Violeae, tribe 71
 Viscaria 44
 Visceae, tribe 38
 Viscoideae, subfamily 38
 Viscum 38

 Wahlenbergia 103
 Wahlenbergiinae, subtribe 103
 Wolffia 23
 Wolfoideae, subfamily 23
 Woodsia 5
 Woodsieae, tribe 5

 Xanthium 105

 Zannichellia 12 1
 Zannichellieae, tribe 12
 Zostera 12
 Zostereae, tribe 12

SELECTION FROM THE GENERAL CATALOGUE
OF BOOKS PUBLISHED BY
THE CAMBRIDGE UNIVERSITY PRESS

The Genus Iris. By WILLIAM RICKATSON DYKES. With 47 coloured drawings by F. H. ROUND, 1 coloured plate of seeds by Miss R. M. CARDEW, and 30 line drawings by C. W. JOHNSON. Demy folio. £6 6s net.

Types of British Vegetation. By members of the Central Committee for the Survey and Study of British Vegetation. Edited by A. G. TANSLEY, M.A., F.L.S., University Lecturer on Botany in the University of Cambridge. Crown 8vo. With 36 plates and 21 figures. 6s net.

The Vegetation of the Peak District. By C. E. MOSS, B.A., D.Sc., F.R.G.S., F.L.S., Curator of the Herbarium, Cambridge. Demy 8vo. With 36 illustrations and 2 coloured maps. 12s net.

A Manual and Dictionary of the Flowering Plants and Ferns. By J. C. WILLIS, M.A., Sc.D. Third edition. Crown 8vo. 10s 6d Cambridge Biological Series.

The Classification of Flowering Plants. By ALFRED BARTON RENDLE, M.A., D.Sc., F.L.S., Keeper of the Department of Botany, British Museum. Vol. I. Gymnosperms and Monocotyledons. Demy 8vo. With 187 illustrations. 10s 6d net. Cambridge Biological Series.

Grasses : a Handbook for use in the Field and Laboratory. By H. MARSHALL WARD, Sc.D., F.R.S. Crown 8vo. With 81 figures. 6s Cambridge Biological Series.

Trees : a Handbook of Forest Botany for the Woodlands and the Laboratory. By H. MARSHALL WARD, Sc.D., F.R.S. Vol. I. Buds and Twigs. Vol. II. Leaves. Vol. III. Flowers and Inflorescences. Vol. IV. Fruits. Vol. V. Form and Habit, with an appendix on Seedlings. Crown 8vo. With numerous illustrations. 4s 6d net each volume. Price for the set of five volumes, 20s net. Cambridge Biological Series.

The Production and Utilization of Pine Timber in Great Britain. Part I, Production. No. 1, Sample Plots of Scots Pine at Woburn. By E. R. BURDON, M.A., Investigator in Timber, and A. P. LONG, B.A., Assistant Investigator. Demy 8vo. Paper covers. 1s 6d net. University of Cambridge School of Forestry Bulletins, No. 1.

British Rust Fungi (Uredinales). Their Biology and Classification. By W. B. GROVE, M.A. Demy 8vo. With 290 figures. 14s net.

Continued overleaf

Rubber and Rubber Planting. By R. H. LOCK, Sc.D., Inspector H.M. Board of Agriculture and Fisheries, sometime Fellow of Gonville and Caius College, Cambridge. Crown 8vo. With 10 plates and 22 figures. 5s net.

The Fertility of the Soil. By E. J. RUSSELL, D.Sc. Royal 16mo. With 9 illustrations. Cloth, 1s net; leather, 2s 6d net. Cambridge Manuals Series.

Mendel's Principles of Heredity. By W. BATESON, M.A., F.R.S., V.M.H., Director of the John Innes Horticultural Institution. Third impression with additions. Royal 8vo. With 3 portraits, 6 coloured plates and 38 figures. 12s net.

Heredity in the Light of Recent Research. By L. DONCASTER, Sc.D. Royal 16mo. With 12 figures. Cloth, 1s net; leather, 2s 6d net. Cambridge Manuals Series.

Herbals. Their Origin and Evolution. A Chapter in the History of Botany, 1470—1760. By AGNES ARBER, D.Sc., F.L.S., Fellow of Newnham College, Cambridge, and of University College, London. Royal 8vo. 10s 6d net.

Makers of British Botany. A collection of Biographies by living botanists, edited by F. W. OLIVER. With frontispiece and 26 illustrations. Demy 8vo. 9s net.

CAMBRIDGE NATURE STUDY SERIES

General Editor: HUGH RICHARDSON, M.A.

Lessons on Soil. By E. J. RUSSELL, D.Sc. Large crown 8vo. With 58 illustrations. 1s 6d net.

The Gateways of Knowledge. An Introduction to the study of the Senses. By J. A. DELL, M.Sc. (Vict.). Large crown 8vo. With 52 illustrations. 2s 6d net.

The Story of Our Trees. In twenty-four lessons. By MARGARET M. GREGSON, B.A. Large crown 8vo. With 75 illustrations. 2s 6d net.

Weeds. Simple Lessons for Children. By R. L. PRAEGER. Large crown 8vo. With 3 plates and 45 illustrations. 1s 6d net.

Cambridge University Press

Fetter Lane, London

C. F. Clay, Manager

